For Research Use Only

CoraLite®555-conjugated Phospho-AKT (Ser473) Recombinant antibody

www.ptglab.com

Catalog Number: CL555-80455

Basic Information

Catalog Number: GenBank Accession Number:

CL555-80455 NM 005163 GeneID (NCBI):

100ul, Concentration: 1000 ug/ml by 207

Nanodrop: **UNIPROT ID:** Source: P31749

Isotype: v-akt murine thymoma viral IgG oncogene homolog 1

> Observed MW: 58 kDa

Full Name:

Purification Method:

Protein A purification

CloneNo.: 2E17

Excitation/Emission maxima

wavelengths: 557 nm / 570 nm

Applications

Tested Applications:

FC (Intra)

Rabbit

Species Specificity:

human

Background Information

AKT is a serine/threonine kinase and it participates in the key role of the PI3K signaling pathway. Phosphatidylinositol-3 kinase (PI3K) is the key regulator of AKT activation. The recruitment of inactive AKT protein to PIP3-rich areas of the plasma membrane results in a conformational change that exposes the activation loop of AKT. AKT's activating kinase, phosphoinositide-dependent protein kinase (PDK1), is also recruited to PIP3 microdomains. PDK1 phosphorylates AKT on threonine 308 (Thr308) of the exposed activation loop, activating AKT and leading to a second phosphorylation of AKT at serine 473 (Ser473) by a kinase presumed to be mTORC2 that $further potentiates \ kinase \ activity. \ Active \ AKT \ will \ phosphory late \ various \ downstream \ protein \ targets \ that \ control$ cell growth and translational control and act to suppress apoptosis. (PMID: 31594388, PMID: 30808672)

Storage

Storage


Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

in USA), or 1(312) 455-8498 (outside USA)

Selected Validation Data



1X10^6 HEK-293 cells untreated (dashed lines) or treated with Calyculin A (red) were intracellularly stained with 0.13 ug Coralite®555 Anti-Human Phospho-AKT (Ser473) (CL555-80455, Clone:2E17), or 0.13 ug Control Antibody (blue). Cells were fixed with 4% PFA and permeabilized with 80% MeOH.