For Research Use Only

## CoraLite®555-conjugated Phospho-AKT1 (Ser473) Recombinant antibody

Catalog Number: CL555-80462



**Basic Information** 

Catalog Number: GenBank Accession Number:

100ul , Concentration: 1000 ug/ml by  $\,$  207  $\,$ 

Nanodrop; UNIPROT ID: Source: P31749

Rabbit Full Name:

Isotype: v-akt murine thymoma viral IgG oncogene homolog 1

Observed MW: 56-62 kDa Purification Method:

Protein A purification

CloneNo.: 2M10

Recommended Dilutions: IF/ICC 1:50-1:500

Excitation/Emission maxima

wavelengths: 557 nm / 570 nm

**Applications** 

Tested Applications: IF/ICC, FC (Intra)

Species Specificity: human, mouse

Positive Controls:

IF/ICC: Calyculin A treated HeLa cells,

## **Background Information**

AKT is a serine/threonine kinase and it participates in the key role of the PI3K signaling pathway. Phosphatidylinositol-3 kinase (PI3K) is the key regulator of AKT activation. The recruitment of inactive AKT protein to PIP3-rich areas of the plasma membrane results in a conformational change that exposes the activation loop of AKT. AKT's activating kinase, phosphoinositide-dependent protein kinase (PDK1), is also recruited to PIP3 microdomains. PDK1 phosphorylates AKT on threonine 308 (Thr308) of the exposed activation loop, activating AKT and leading to a second phosphorylation of AKT at serine 473 (Ser473) by a kinase presumed to be mTORC2 that further potentiates kinase activity. Active AKT will phosphorylate various downstream protein targets that control cell growth and translational control and act to suppress apoptosis. (PMID: 31594388, PMID: 30808672). 80462-1-RR specifically recognizes AKT1 phosphorylated at Ser473.

Storage

Storage:


Store at -20 °C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer


PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

## **Selected Validation Data**



1X10^6 NIH/3T3 cells untreated (dashed line) or treated with Calyculin A (red) were intracellularly stained with 0.25 ug CoraLite®555 Anti-Human Phospho-AlKT1 (Ser473) (CL555-80462, Clone:2M10), or 0.25 ug Control Antibody (Blue). Cells were fixed with 4% PFA and permeabilized with 90% MeOH.



Immunofluorescent analysis of (4% PFA) fixed Calyculin A treated Hela cells using Coralite®555 Phospho-AKT1 (Ser473) antibody (CL555-80462, Clone: 2M10 ) at dilution of 1:200.