For Research Use Only

CoraLite®594-conjugated PAX8 Polyclonal antibody

Catalog Number: CL594-10336



Basic Information

Catalog Number: GenBank Accession Number:

Observed MW:

CL594-10336 BC001060 GeneID (NCBI):

100ul, Concentration: 1000 ug/ml by 7849 Nanodrop:

UNIPROT ID: Q06710 Rabbit Full Name: Isotype: paired box 8 IgG Calculated MW: Immunogen Catalog Number: 48 kDa AG0306

Purification Method: Antigen affinity purification Recommended Dilutions: IF/ICC 1:50-1:500

Excitation/Emission maxima wavelengths:

588 nm / 604 nm

Applications

Tested Applications: IF/ICC, FC (Intra) Species Specificity:

human, mouse, rat

Positive Controls: IF/ICC: SKOV-3 cells,

Background Information

PAX8 is a member of the paired box (PAX) family of transcription factors, typically containing a paired box domain $and\ a\ paired-type\ homeodomain.\ It\ is\ expressed\ during\ organogenesis\ of\ the\ thyroid\ gland,\ kidney\ and\ Mullerian$ system. It is thought to regulate the expression of Wilms tumor suppressor (WT1) gene and mutations in PAX8 have been associated with Wilms tumor cells, thyroid and ovarian carcinomas. PAX8 is a useful marker in distinguishing ovarian carcinomas from mammary carcinomas (PMID: 18724243). PAX8 is expressed in a high percentage of ovarian serous, endometrioid, and clear cell carcinomas, but only rarely in primary ovarian mucinous adenocarcinomas.

This antibody can detect all 5 isoforms of PAX8 (31 kDa, 34 kDa, 41 kDa, 43 kDa and 48 kDa). A 58-60 kDa form has also been reported (PMID:15650356, 1049952). Proteintech's 10336-1-AP antibody has cross-reactivities with the other PAX family members.

Storage

Storage:


Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.


Aliquoting is unnecessary for -20°C storage

in USA), or 1(312) 455-8498 (outside USA)

Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed SKOV-3 cells using CoraLite®594 PAX8 antibody (CL594-10336) at dilution of 1:200.



1X10^6 SKOV-3 cells were intracellularly stained with 0.2 ug Coralite®594 Anti-Human PAX8 (CL594-10336) (red), or 0.2 ug Control Antibody. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).