For Research Use Only

## CoraLite®594-conjugated PAX8 Polyclonal antibody



Catalog Number: CL594-10336

**Basic Information** 

Catalog Number: GenBank Accession Number: CL594-10336

BC001060

GeneID (NCBI):

100ul, Concentration: 1000 µg/ml by 7849

Full Name:

Source: paired box 8 Rabbit Calculated MW:

Isotype: 48 kDa IgG Observed MW:

Immunogen Catalog Number:

AG0306

**Purification Method:** Antigen affinity purification Recommended Dilutions:

IF 1:50-1:500

Excitation/Emission maxima

wavelengths: 593 nm / 614 nm

**Applications** 

**Tested Applications:** 

FC (Intra), IF

Species Specificity: human, mouse, rat

**Positive Controls:** 

IF: SKOV-3 cells,

## **Background Information**

PAX8 is a member of the paired box (PAX) family of transcription factors, typically containing a paired box domain and a paired-type homeodomain. It is expressed during organogenesis of the thyroid gland, kidney and Mullerian system. It is thought to regulate the expression of Wilms tumor suppressor (WT1) gene and mutations in PAX8 have been associated with Wilms tumor cells, thyroid and ovarian carcinomas. PAX8 is a useful marker in distinguishing ovarian carcinomas from mammary carcinomas (PMID: 18724243). PAX8 is expressed in a high percentage of ovarian serous, endometrioid, and clear cell carcinomas, but only rarely in primary ovarian mucinous adenocarcinomas.

This antibody can detect all 5 isoforms of PAX8 (31 kDa, 34 kDa, 41 kDa, 43 kDa and 48 kDa). A 58-60 kDa form has also been reported (PMID:15650356, 1049952). Proteintech's 10336-1-AP antibody has cross-reactivities with the other PAX family members.

Storage

Store at -20°C. Avoid exposure to light.


Storage Buffer:

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.


Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA

## Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed SKOV-3 cells using CoraLite®594 PAX8 antibody (CL594-10336) at dilution of 1:200.



1X10^6 SKOV-3 cells were intracellularly stained with 0.2 ug Coralite®594 Anti-Human PAX8 (CL594-10336) (red), or 0.2 ug Control Antibody. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).