For Research Use Only

## CoraLite®594-conjugated Emerin Polyclonal antibody

Catalog Number:CL594-10351 Featured Product



**Basic Information** 

Catalog Number: GenBank Accession Number:

CL594-10351 BC000738 GeneID (NCBI):

100ul, Concentration: 1000 ug/ml by 2010

Nanodrop: **UNIPROT ID:** P50402 Rabbit Full Name:

Isotype: emerin

IgG Calculated MW: 34 kDa

Immunogen Catalog Number: AG0236

Observed MW: 35 kDa

**Purification Method:** Antigen affinity purification

Recommended Dilutions: IF/ICC 1:50-1:500

Excitation/Emission maxima

wavelengths: 588 nm / 604 nm

**Applications** 

**Tested Applications:** IF/ICC, FC (Intra) Species Specificity:

human, mouse

Positive Controls: IF/ICC: HeLa cells,

**Background Information** 

Emerin (Emery-Dreifuss muscular dystrophy) (EMD or EDMD), a serine-rich nuclear membrane protein, is a member of the nuclear lamina-associated protein family. EMD may mediate membrane anchorage to the cytoskeleton by stabilizing and promoting the formation of a nuclear actin cortical network. Defects in EMD gene are the cause of Emery-Dreifuss muscular dystrophy type 1 (EDMD1), a degenerative myopathy characterized by weakness and atrophy of muscle without involvement of the nervous system, early contractures of the elbows Achilles tendons and spine, and cardiomyopathy associated with cardiac conduction defects. EMD inhibits beta-catenin activity by preventing its accumulation in the nucleus and is involved in HIV-1 infection.

Storage

Storage:

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.


Storage Buffer:

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.


Aliquoting is unnecessary for -20°C storage

in USA), or 1(312) 455-8498 (outside USA)

## Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using CoraLite®594 Emerin antibody (CL594-10351) at dilution of 1:100.



1X10^6 HeLa cells were intracellularly stained with 0.4 ug CoraLite® 594 Anti-Human Emerin (CL594-10351) (red), or 0.4 ug Control Antibody. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).