For Research Use Only

CoraLite®594-conjugated P21 Polyclonal antibody

Catalog Number:CL594-10355 Featured Product



Purification Method:

Basic Information

Catalog Number: GenBank Accession Number:

CL594-10355 BC000275 Antigen affinity purification
Size: GeneID (NCBI): Recommended Dilutions:

100ul , Concentration: 1000 ug/ml by 1026 IF/ICC 1:50-1:500

Nanodrop;UNIPROT ID:Excitation/Emission maximaSource:P38936wavelengths:RabbitFull Name:588 nm / 604 nm

Isotype: cyclin-dependent kinase inhibitor 1A

IgG (p21, Cip1)
Immunogen Catalog Number: Calculated MW:
AG0368 18 kDa

Observed MW: 21 kDa

Applications

Tested Applications: Positive Controls: IF/ICC, FC (Intra) IF/ICC : MCF-7 cells,

Species Specificity: human

Background Information

CDKN1A (p21, CIP1, WAF1) is a cyclin-dependent kinase inhibitor. CDKN1A binds to and inhibits the activity of cyclin-CDK2 or -CDK4 complexes, and thus functions as a regulator of cell cycle progression at the G1 phase. The expression of CDKN1A is induced by wild-type but not mutant p53 protein, through which CDKN1A mediates the p53-dependent cell cycle G1 phase arrest in response to a variety of stress stimuli. CDKN1A can interact with proliferating cell nuclear antigen (PCNA), and plays a regulatory role in S phase DNA replication and DNA damage repair. CDKN1A was reported to be specifically cleaved by CASP3-like caspases, which thus leads to a dramatic activation of CDK2, and may be instrumental in the execution of apoptosis following caspase activation. Two alternatively spliced variants, which encode an identical protein, have been reported.

Storage


Storage:

Store at -20°C. Avoid exposure to light. Stable for one year after shipment. Storage Buffer:


PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed MCF-7 cells using Coralite®594 P21 antibody (CL594-10355) at dilution of 1:100.



1X10^6 MCF-7 cells were intracellularly stained with 0.8 ug CoraLite®594 Anti-Human P21 (CL594-10355) (red), or 0.8 ug CoraLite®594-conjugated Rabbit IgG control Rabbit PolyAb (CL594-30000, Clone:) (blue). Cells were fixed and permeabilized with True-Nuclear Transcription Factor Buffer Set.