## For Research Use Only

## CoraLite®594-conjugated MMP-9 (Nterminal) Polyclonal antibody

www.ptglab.com

Catalog Number: CL594-10375

Featured Product

1 Publications

**Basic Information** 

Catalog Number:

GenBank Accession Number:

Antigen affinity purification

CL594-10375

BC006093 GeneID (NCBI):

Recommended Dilutions:

100ul, Concentration: 1000 ug/ml by 4318

IF/ICC 1:50-1:500

**Purification Method:** 

588 nm / 604 nm

Nanodrop:

**UNIPROT ID:** P14780

Excitation/Emission maxima wavelengths:

Rabbit

Full Name:

Isotype IgG

matrix metallopeptidase 9 (gelatinase B, 92kDa gelatinase, 92kDa type IV collagenase)

Immunogen Catalog Number: AG0552

Calculated MW:

707 aa, 78 kDa

Observed MW:

67 kDa, 92 kDa

**Applications** 

**Tested Applications:** 

IF/ICC, FC (Intra)

**Cited Applications:** 

IF

IF/ICC: HepG2 cells,

**Positive Controls:** 

Species Specificity: human, mouse, rat

**Cited Species:** 

human

**Background Information** 

Proteins of the matrix metalloproteinase (MMP) family are involved in the breakdown of extracellular matrix in normal physiological processes, such as embryonic development, reproduction, tissue remodeling, and disease processes, such as arthritis or metastasis. Most MMP's are secreted as inactive proproteins which are activated when cleaved by extracellular proteinases. Matrix metalloproteinase 9 (gelatinase B, 92kDa gelatinase, 92kDa type IV collagenase) (MMP9, synonyms: GELB, CLG4B) degrades collagens type IV and V. Studies in rhesus monkeys suggest that MMP9 is involved in IL-8-induced mobilization hematopoietic progenitor cells from bone marrow, and murine studies suggest a role in tumor-associated tissue remodeling. The pro-MMP9 is 92 kDa, and it can be detected a processed form of 68 kDa. This protein can exist as a dimer of 180 kDa (PMID:7492685).

**Notable Publications** 

Author Pubmed ID Application Journal Qian Jiang 37444485 Cancers (Basel)

Storage


Storage:

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.


PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

## Selected Validation Data



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using CoraLite®594 MMP9 (N-terminal) antibody (CL594-10375) at dilution of 1:200.



1X10^6 HeLa cells were intracellularly stained with 0.4 ug CoraLite® 594 Anti-Human MMP9 (N-terminal) (CL594-10375) (red), or 0.4 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).