For Research Use Only

## CoraLite®594-conjugated TFEB Polyclonal antibody

Catalog Number:CL594-13372

Featured Product






Basic Information	Catalog Number: CL594-13372	GenBank Accession Number: BC032448	Purification Method: Antigen affinity purification	
	Size: 100ul , Concentration: 1000 ug/ml by	GenelD (NCBI): 7942	Recommended Dilutions: IF/ICC 1:50-1:500	
	Nanodrop; Source: Rabbit Isotype: IgG Immunogen Catalog Number: AG4194	UNIPROT ID: P19484 Full Name: transcription factor EB Calculated MW: 476 aa, 53 kDa Observed MW: 65-70 kDa	Excitation/Emission maxima wavelengths: 588 nm / 604 nm	
Applications	Tested Applications: IF/ICC		Controls: JepG2 cells,	
	Cited Applications: IF Species Specificity: human, mouse, rat			
Background Information	Transcription factor EB(TFEB) is one of the transcription factors that contain a basic HLH-ZIP domain(for DNA binding ) and HLH and ZIP domains( for loigomerization). TFEB specifically recognizes and binds to E-box sequences, which required dimerization with itself or another MiT/TFE family member. It plays a role in T-cell-dependent antibody responses to activated CD4+ T-cells and thymus-dependent humoral immunity once associated with TFE3. It also activates lysosomal gene by binding the CLEAR-box sequence in the regulatory region. It involves in the signal transduction processes required for normal vascularization of the placenta. This is a rabbit polyclonal antibody raised against the full length of human TFEB. The calcualted molecular weight of TFEB is 53 kDa, but modified TFEB is about 65-70 kDa (PMID: 12837690). TFEB mainly present in the cytoplasm (PubMed:23434374). Under aberrant lysosomal storage conditions, it translocates from the cytoplasm to the nucleus (PubMed:23434374). Fisetin increases the levels of TFEB localized in nuclei (PMID: 27112200).			
Notable Publications	Author Pr	ibmed ID Journal	Application	
	Margot R F Reijnders 37	057673 Genet Med	IF	
Storage	Storage: Store at -20°C. Avoid exposure to lig Storage Buffer: PBS with 50% Glycerol, 0.05% Procl Aliquoting is unnecessary for -20°C	in300, 0.5% BSA, pH 7.3.	pment.	


For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free<br/>in USA), or 1(312) 455-8498 (outside USA)E: proteintech@ptglab.comW: ptglab.comW: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data



Immunofluorescent analysis of (-20°C Methanol) fixed HepG2 cells using Coralite®594 TFEB antibody (CL594-13372) at dilution of 1:200.



Immunofluorescent analysis of (-20°C Methanol) fixed HepG2 cells using CoraLite®594 TFEB antibody (CL594-13372) at dilution of 1:600.