For Research Use Only

CoraLite®594-conjugated Fibronectin Polyclonal antibody

Proteintech®
Antibodies | ELISA kits | Proteins
www.ptglab.com

Purification Method:

IF/ICC 1:50-1:500

wavelengths: 588 nm / 604 nm

Antigen affinity purification

Excitation/Emission maxima

Recommended Dilutions:

Catalog Number: CL594-15613

Featured Product

Basic Information

Catalog Number: GenBank Accession Number:

CL594-15613 BC005858 Size: GeneID (NCBI):

100ul , Concentration: 1000 ug/ml by 2335

Nanodrop; UNIPROT ID:
Source: P02751
Rabbit Full Name:

Isotype:fibronectin 1IgGCalculated MW:Immunogen Catalog Number:2386 aa, 263 kDa

AG8004

Applications

Tested Applications: IF/ICC, FC (Intra)

Species Specificity: human, mouse

Positive Controls:

IF/ICC: NIH/3T3 cells,

Background Information

Fibronectin 1 (FN1) is a high molecular weight glycoprotein which exists in both a soluble form in plasma (plasma FN1) and other body fluids and an insoluble form in the extracellular matrix (cellular FN1). Plasma FN1 (dimeric form) is secreted by hepatocytes. Cellular FN (dimeric or cross-linked multimeric forms), made by fibroblasts, epithelial and other cell types, is deposited as fibrils in the extracellular matrix. FN1 binds to cell surfaces through integrins and to various compounds including collagen, fibrin and heparin. It is involved in cell adhesion and migration processes including embryogenesis, wound healing, hemostasis, host defense, and metastasis. (PMID: 7276612; 12244123; 12847088)

Storage

Storage:


Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer


PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Immunofluorescent analysis of (-20°C Methanol) fixed NIH/3T3 cells using CoraLite®594 Fibronectin antibody (CL594-15613) at dilution of 1:200.



1X10^6 NIH/3T3 cells were intracellularly stained with 0.4 ug CoraLite®594 Anti-Human Fibronectin (CL594-15613) (red), or 0.4 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).