

For Research Use Only

CoraLite®594-conjugated MAP2 Polyclonal antibody

Catalog Number: CL594-17490

2 Publications



Basic Information

Catalog Number:

CL594-17490

Size:

100ul , Concentration: 1000 ug/ml by
Nanodrop;

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG11580

GenBank Accession Number:

BC038857

GeneID (NCBI):

4133

UNIPROT ID:

P11137

Full Name:

microtubule-associated protein 2

Calculated MW:

200 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

IF-P: 1:50-1:500

FC (Intra): 0.40 ug per 10⁶ cells in a
100 µl suspension

Excitation/Emission maxima
wavelengths:

588 nm / 604 nm

Applications

Tested Applications:

IF-P, FC (Intra)

Cited Applications:

IF

Species Specificity:

human, mouse, rat

Cited Species:

mouse, rat

Positive Controls:

IF-P: rat brain tissue,

FC (Intra): Neuro-2a cells,

Background Information

MAP2 (microtubule-associated protein 2) is a cytoskeleton protein abundant in brain and has important role in neuronal morphogenesis. Multiple high MW and low MW MAP2 isoforms are expressed within proximal segment of axons, dendrites, and cell bodies. The expression of MAP2 is regulated in both a tissue- and developmentally specific manner. The 280 kDa MAP2B is present throughout rat brain development, and the slightly larger MAP2A appears first during the end of the second week of postnatal life. MAP2C, composed of several bands of about 70 kDa, is present during early brain development, and largely disappears from the mature brain except for the retina, olfactory bulb, and cerebellum. In addition, some isoforms with lower MW around 50-60 kDa also exist. MAP2 antibodies have been widely used to mark the neuron or dendrite formation. This antibody can recognize both high MW and low MW isoforms of MAP2.

Notable Publications

Author	Pubmed ID	Journal	Application
Jian Lu	40041886	Acta Pharm Sin B	
Annette Vaglio-Garro	38785950	Biomolecules	IF

Storage

Storage:

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer:

PBS with 50% glycerol, 0.05% Proclin300, 0.5% BSA, pH7.3

Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed rat brain tissue using CoraLite®594 MAP2 antibody (CL594-17490) at dilution of 1:200, CoraLite®488 GFAP antibody (CL488-16825, green). DAPI (blue).



Immunofluorescent analysis of (4% PFA) fixed rat brain tissue using CoraLite®594 MAP2 antibody (CL594-17490) at dilution of 1:200, CoraLite®488 GFAP antibody (CL488-16825, green). DAPI (blue).



1X10⁶ Neuro-2a cells were intracellularly stained with 0.4 ug CoraLite®594 Anti-Human MAP2 (CL594-17490) (red), or 0.4 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).