### For Research Use Only

# CoraLite®594-conjugated VEGFA Polyclonal antibody



**Purification Method:** 

Catalog Number: CL594-19003

Featured Product

**Basic Information** 

Catalog Number: GenBank Accession Number:

CL594-19003 BC065522 Antigen affinity purification GeneID (NCBI): Recommended Dilutions: IF/ICC 1:50-1:500

100ul, Concentration: 1000 ug/ml by 7422 Nanodrop: **UNIPROT ID:** Excitation/Emission maxima

P15692 wavelengths: 588 nm / 604 nm Rabbit Full Name:

Isotype: vascular endothelial growth factor A

IgG Calculated MW: Immunogen Catalog Number: 412 aa, 46 kDa

AG13500

**Applications** 

**Tested Applications:** Positive Controls: IF/ICC, FC (Intra) IF/ICC: HeLa cells,

Species Specificity: human, mouse

#### **Background Information**

VEGFA, also named as VEGF or VPF, belongs to the PDGF/VEGF growth factor family. It is a growth factor active in angiogenesis, vasculogenesis and endothelial cell growth. VEGFA induces endothelial cell proliferation, promotes cell migration, inhibits apoptosis and induces permeabilization of blood vessels. It binds to the FLT1/VEGFR1 and KDR/VEGFR2 receptors, heparan sulfate and heparin. Defects in VEGFA are associated with microvascular complications of diabetes type 1 (MVCD1). VEGFA has 17 isoforms with MW from 16 to 45 kDa. Some isoforms have homodimer forms (e.g.; VEGFA189 38kDa or VEFGA110 34kDa). VEGF-A exists in at least seven homodimeric  $is of orms. The \,monomers \,consist \,of \,121,\,145,\,148,\,165,\,183,\,189,\,or \,206 \,amino \,acids \,(PMID:15602010\,). \,This \,antibody \,acids \,(PMID:15602010\,).$ can recognize all VEGFA isoforms.

Storage

#### Storage:


Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.


Aliquoting is unnecessary for -20°C storage

in USA), or 1(312) 455-8498 (outside USA)

## Selected Validation Data



Immunofluorescent analysis of (-20°C Ethanol) fixed HeLa cells using CoraLite®594 VEGF antibody (CL594-19003) at dilution of 1:100.



1X10^6 NIH/3T3 cells were intracellularly stained with 0.4 ug Coralite®594 Anti-Human VEGF (CL594-19003) (red), or 0.4 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).