For Research Use Only

CoraLite®594-conjugated NLRP3 Polyclonal antibody



Catalog Number: CL594-19771

Featured Product

Basic Information

Catalog Number: GenBank Accession Number:

CL594-19771 NM_001127461 GeneID (NCBI): 100ul , Concentration: 1000 $\mu g/ml$ by 114548

Nanodrop: **UNIPROT ID:** Source: Q96P20 Rabbit Full Name:

Isotype: NLR family, pyrin domain containing

IgG

Calculated MW: 118 kDa Observed MW: 110 kDa

Purification Method:

Antigen affinity purification Excitation/Emission maxima

wavelengths: 588 nm / 604 nm

Applications

Tested Applications:

FC (Intra)

Species Specificity:

Background Information

NALP3, also named as C1orf7, CIAS1 and PYPAF1, belongs to the NLRP family. NALP3 may function as an inducer of $apoptosis. It interacts selectively with ASC and this complex \ may function as an upstream \ activator \ of \ NF-kappa-B$ signaling.NALP3 inhibits TNF-alpha induced activation and nuclear translocation of RELA/NF-KB p65. Also inhibits transcriptional activity of RELA. NALP3 activates caspase-1 in response to a number of triggers including bacterial or viral infection which leads to processing and release of IL1B and IL18. Defects in NLRP3 are the cause of familial cold autoinflammatory syndrome type 1 (FCAS1) which also known as familial cold urticaria. Defects in NLRP3 are a $cause of \, Muckle-Wells \, syndrome \, (MWS) \, which \, is \, urticaria-deafness-amyloidos is \, syndrome. \, Defects \, in \, NLRP3 \, are \, the \, in \, NLRP3 \, are \, the \, in \, NLRP3 \, and \, in \, NLRP3 \, are \, the \, in \, NLRP3 \, and \, in \, NLRP3 \,$ cause of chronic infantile neurologic cutaneous and articular syndrome (CINCA) which also known as neonatal onset multisystem inflammatory disease (NOMID). The antibody recognizes the C-term of NALP3.

Storage


Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

in USA), or 1(312) 455-8498 (outside USA)

Selected Validation Data



1X10^6 THP-1 cells were intracellularly stained with 0.4 ug CoraLite® 594 Anti-Human NLRP3 (CL594-19771) (red), or 0.4 ug Isotype Control. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).