For Research Use Only

## CoraLite®594-conjugated NFATC2 Polyclonal antibody



**Purification Method:** 

Catalog Number: CL594-22023

Featured Product

**Basic Information** 

Catalog Number: GenBank Accession Number:

CL594-22023 BC136418 Antigen affinity purification GeneID (NCBI): Recommended Dilutions: 100ul, Concentration: 1000 ug/ml by 4773 IF/ICC 1:50-1:500

Nanodrop: **UNIPROT ID:** Excitation/Emission maxima

Q13469 wavelengths: 588 nm / 604 nm Rabbit Full Name:

Isotype nuclear factor of activated T-cells, IgG cytoplasmic, calcineurin-dependent 2

Immunogen Catalog Number: Calculated MW: AG16849 925 aa. 100 kDa

Observed MW: 135 kDa

**Applications** 

**Tested Applications:** Positive Controls: IF/ICC, FC (Intra) IF/ICC: U-251 cells,

Species Specificity: human, mouse

## **Background Information**

Nuclear factor of activated T-cells, cytoplasmic 2 (NFATC2), also named NFAT1, or NFATP, is a 925 amino acid protein, which is expressed in thymus, spleen, heart, testis, brain, placenta, muscle and pancreas. Cytoplasmic for the phosphorylated form and nuclear after activation that is controlled by calcineurin-mediated dephosphorylation. Rapid nuclear exit of NFATC is thought to be one mechanism by which cells distinguish between sustained and transient calcium signals. The subcellular localization of NFATC plays a key role in the regulation of gene transcription. NFATC2 plays a role in the inducible expression of cytokine genes in T-cells, especially in the induction of the IL-2, IL-3, IL-4, TNF-alpha or GM-CSF. NFATC2 promotes invasive migration through the activation of GPC6 expression and WNT5A signaling pathway. The calculated molecular weight of NFATC2 is about 97-100 kDa, but the modified NFATC2 protein is about 135 kDa.

Storage

## Storage:


Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.


Aliquoting is unnecessary for -20°C storage

in USA), or 1(312) 455-8498 (outside USA)

## Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed U-251 cells using CoraLite®594 NFATC2 antibody (CL594-22023) at dilution of 1:200.



1X10^6 Ramos cells were intracellularly stained with 0.4 ug Coralite®594 Anti-Human NFATC2 (CL594-22023) (red), or 0.4 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).