For Research Use Only

## CoraLite®594-conjugated GRP94 Monoclonal antibody



Catalog Number: CL594-60012

Basic Information

Catalog Number: GenBank Accession Number: Purification Method: Protein G purification

CL594-60012

GenBank Accession Number: Purification Method: Protein G purification

Size: GeneID (NCBI): CloneNo.: 100ul , Concentration: 1000 µg/ml by 7184 4G7C7

anodrop; Full Name: Recommended Dilutions:

Source:heat shock protein 90kDa betaWB 1:500-1:1000Mouse(Grp94), member 1IF 1:50-1:500

 Isotype:
 Calculated MW:
 Excitation/Emission maxima

 IgG1
 96 kDa
 wavelengths:

Immunogen Catalog Number: Observed MW: 588 nm / 604 nm AG1439 95 kDa

Applications Tested Applications: Positive Controls:

FC (Intra), IF, WB WB: HeLa cells, HEK-293 cells, Jurkat cells, HepG2

Species Specificity:

Human IF : HepG2 cells,

## **Background Information**

HSP90 proteins are highly conserved molecular chaperones, which normally associate with other cochaperones and play important roles in folding newly synthesized proteins or stabilizing and refolding denatured proteins after stress. HSP90B1 (GP96 or GRP94) is an endoplasmic reticulum paralogue of the cytosolic HSP90. As a major ER chaperone to mediate the UPR and a master chaperone for Toll-like receptors (TLRs), HSP90b1 chaperones peptides to MHC class I molecules of dendritic cells and other antigen-presenting cells, as well as facilitating the assembly of immunoglobulin. The protein is also involved in many other bio-processes. This antibody was generated against the N-terminal region of full-length HSP90b1.

## Storage

Storage:


Store at -20°C. Avoid exposure to light. Stable for one year after shipment. Storage Buffer:

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


## **Selected Validation Data**



Various lysates were subjected to SDS PAGE followed by western blot with CL594-60012 (GRP94 antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using Coralite®594-conjugated GRP94 antibody (CL594-60012, Clone: 4G7C7) at dilution of 1:100.



1X10^6 HeLa cells were intracellularly stained with 0.4 ug CoraLite®594 Anti-Human GRP94 (CL594-60012, Clone:4G7C7) (red), or 0.4 ug Mouse IgG1 Isotype Control (CL594-66360, Clone: T1F8D3F10) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).