For Research Use Only

CoraLite®594-conjugated Stanniocalcin 2 Monoclonal antibody

Catalog Number: CL594-60063

Featured Product



wavelengths:

588 nm / 604 nm

Basic Information

Catalog Number: GenBank Accession Number: Purification Method: Protein G purification

CL594-60063 BC000658 Protein G purification

Size: GeneID (NCBI): CloneNo.: 100ul , Concentration: 1000 ug/ml by 8614 4H1E7

Nanodrop; UNIPROT ID: Recommended Dilutions:

Source: 076061 IF/ICC: 1:50-1:500

Mouse Full Name: FC (Intra): 0.50 ug per 10^6 cells in a

Isotype: stanniocalcin 2 100 μl suspension
IgG1 Excitation/Emission maxima

IgG1 Calculated MW:
Immunogen Catalog Number: 33 kDa

AG0359 Observed MW:

33 kDa

Applications Tested Applications: Positive Controls:

IF/ICC, FC (Intra)

Species Specificity:
human, mouse

IF/ICC : HeLa cells,
FC (Intra) : HeLa cells,

Background Information

Stanniocalcin (STC) is a family of secreted glycoprotein hormones that originally discovered in the corpuscles of Stannius, an endocrine gland of fish. STC1 and STC2, two homologues of STC family, are reported to involve in calcium and phosphate homeostasis. It is expressed in a wide variety of tissues such as kidney, spleen, heart, and pancreas. The protein may play a role in the regulation of renal and intestinal calcium and phosphate transport, cell metabolism, or cellular calcium/phosphate homeostasis. STC2 overexpression could promote tumor cell proliferation, invasion and metastasis in prostate cancer, ovarian cancer or neuroblastoma. STC2 is also vital for cytoprotective properties when exposed to ER stress and hypoxia.

Storage


Storage

Store at -20°C. Avoid exposure to light. Stable for one year after shipment. Storage Buffer:


PBS with 50% glycerol, 0.05% Proclin300, 0.5% BSA, pH7.3

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Immunofluorescent analysis of (-20°C Ethanol) fixed HeLa cells using CL594-60063 (Stanniocalcin 2 antibody) at dilution of 1:100.



1X10^6 HeLa cells were intracellularly stained with 0.5 ug CoraLite® 594 Anti-Human Stanniocalcin 2 (CL594-60063, Clone: AH1E7) (red), or 0.5 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).