

For Research Use Only

CoraLite®594-conjugated Neuropilin 1 Monoclonal antibody



Catalog Number:CL594-60067

Basic Information

Catalog Number: CL594-60067	GenBank Accession Number: BC007737	Purification Method: Protein G purification
Size: 100ul , Concentration: 2000 µg/ml by Nanodrop;	GeneID (NCBI): 8829	CloneNo.: 2H3F6
Source: Mouse	Full Name: neuropilin 1	Excitation/Emission maxima wavelengths: 588 nm / 604 nm
Isotype: IgG1	Calculated MW: 103 kDa	
Immunogen Catalog Number: AG0931	Observed MW: 130 kDa	

Applications

Tested Applications:
FC (Intra)

Species Specificity:
human, pig

Background Information

Neuropilin-1 (NRP1) is a 130-140 kDa transmembrane glycoprotein expressed by endothelial, dendritic, and regulatory T cells, as well as several other normal cell types and malignant tumor cells. NRP1 was first identified as a semaphorin (SEMA) receptor, involved in axonal guidance in embryonic development. NRP1 was also shown to act as a receptor for vascular endothelial growth factor (VEGF) and a promoter of angiogenesis through its interaction with VEGF-A165 (and other VEGFs) and the receptor tyrosine kinase (RTK) VEGF-R2. NRP1 plays versatile roles in angiogenesis, axon guidance, cell survival, migration, and invasion.

Storage

Storage:
Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer:
PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA) E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



1X10⁶ SH-SY5Y cells were intracellularly stained with 0.4 ug CoraLite® 594 Anti-Human Neuropilin 1 (CL594-60067, Clone:2H3F6) (red), or 0.4 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).