For Research Use Only

## CoraLite®594-conjugated STAT3 Monoclonal antibody



**Purification Method:** 

Catalog Number: CL594-60199

Featured Product

**Basic Information** 

Catalog Number: GenBank Accession Number: BC000627

CL594-60199 Protein A purification GeneID (NCBI): CloneNo.:

100ul, Concentration: 1000 ug/ml by 6774 3G2D12

Nanodrop: **UNIPROT ID:** Recommended Dilutions: P40763 IF/ICC 1:50-1:500

Mouse Excitation/Emission maxima Full Name:

Isotype signal transducer and activator of wavelengths: transcription 3 (acute-phase response 588 nm / 604 nm IgG2a

factor) Immunogen Catalog Number:

Calculated MW: AG0360

> 770 aa, 88 kDa Observed MW: 88 kDa

**Applications** 

**Tested Applications:** 

IF/ICC

Species Specificity:

human, mouse, rat

**Positive Controls:** 

IF/ICC: HepG2 cells,

## **Background Information**

Signal transducer and activator of transcription 3 (acute-phase response factor) (STAT3, synonyms: APRF, FLJ 20882, MGC 16063) is a member of the STAT protein family. In response to cytokines and growth factors, STAT family members are phosphorylated by the receptor associated kinases, and then form homo- or heterodimers that translocate to the cell nucleus where they act as transcription activators. STAT3 is activated through phosphorylation in response to various cytokines and growth factors including IFNs, EGF, IL6, HGF, LIF and BMP2. STAT3 mediates the expression of a variety of genes in response to cell stimuli, and thus plays a key role in many cellular processes such as cell growth and apoptosis. The small GTPase Rac1 has been shown to bind and regulate the activity of STAT3. This antibody is a mouse monoclonal antibody raised against residues near the N terminus of human STAT3. STAT3 exists three isoforms and the molecular weight of each isoform respectively is 83 kDa, 87 kDa and 88 kDa.

Storage


Storage:

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

## Selected Validation Data



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using CL594-60199 (STAT3 antibody) at dilution of 1:100.