For Research Use Only

## CoraLite®594 Anti-Mouse IL-12/IL-23 p40 (C17.8)



Catalog Number: CL594-65097

Basic Information Catal

Catalog Number:

CL594-65097

100ug , 500 µg/ml

Source: Rat

Isotype: IgG2a, kappa GenBank Accession Number:

BC103608 GeneID (NCBI):

16160

Full Name: interleukin 12b

Purification Method: Affinity purification

CloneNo.:

Excitation/Emission maxima

wavelengths: 588 nm / 604 nm

**Applications** 

**Tested Applications:** 

FC (Intra)

Species Specificity:

Mouse

**Background Information** 

Interleukin (IL)-12 and IL-23 are heterodimeric cytokines that share a common p40 subunit (PMID: 11114383). IL-12 is composed of the IL-12 p40 subunit linked to the IL-12 p35 subunit, and the heterodimer signals through the IL-12 receptor (IL-12R), which comprises the IL-12R $\beta$ 1 and IL-12R $\beta$ 2 subunits. IL-23 is composed of the IL-23 p19 subunit and the IL-12 p40 (IL-12/23p40) subunit, which signals through IL-23R and IL-12R $\beta$ 1 (PMID: 11114383; 26121196 ). IL-12/IL-23 p40 also exists as a monomer and as a homodimer which can act as a potent IL-12 antagonist (PMID: 8958912; 18783467). IL-12/IL-23 p40 is produced by antigen-presenting cells, such as dendritic cells (DCs), monocytes, macrophages, neutrophils and, to a lesser extent, B cells (PMID: 20476918).

Storage


Storage

Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer:

PBS with 0.09% sodium azide.

## Selected Validation Data



1X10^6 IFN-Y primed Balb/c mouse peritoneal macrophages were unstimulated or stimulated with LPS in the presence of a protein transport inhibitor and then intracellularly stained with 0.5 ug CoraLite®594 Anti-Mouse IL-12/IL-23 p40 (CL594-65097, Clone:C17.8). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).