

For Research Use Only

CoraLite®594 Anti-Mouse TCR beta (H57-597) Armenian Hamster Recombinant Antibody

Catalog Number: CL594-65678



Basic Information

Catalog Number:

CL594-65678

Size:

100ug, 500 ug/ml

Source:

Armenian Hamster

Isotype:

IgG2a

GenBank Accession Number:

GeneID (NCBI):

21577

Full Name:

T-cell receptor beta chain

Purification Method:

Protein A&G purification

CloneNo.:

H57-597

Recommended Dilutions:

FC: 0.25 ug per 10⁶ cells in a 100 µl suspension

Excitation/Emission maxima wavelengths:

588 nm / 604 nm

Applications

Tested Applications:

FC

Species Specificity:

mouse

Positive Controls:

FC : mouse splenocytes,

Background Information

T-cell receptor (TCR) consists of heterodimeric glycoproteins (TCR alpha/beta or gamma/delta) that demonstrate homology with immunoglobulins (PMID: 2144901). TCR beta is a member of the immunoglobulin superfamily and is expressed on alpha/beta TCR-bearing T cells and thymocytes. The TCR heterodimers are noncovalently associated with CD3. The TCR/CD3 complex of T-lymphocytes consists of either a TCR alpha/beta or TCR gamma/delta heterodimer coexpressed at the cell surface with the invariant subunits of CD3 labeled gamma, delta, epsilon, zeta, and eta (PMID: 1826255). The TCR recognizes antigens bound to major histocompatibility complex (MHC) molecules. TCR-mediated peptide-MHC recognition is transmitted to the CD3 complex, leading to the intracellular signal transduction (PMID: 11985657).

Storage

Storage:

Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer:

PBS with 0.09% sodium azide, pH7.3


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



1x10⁶ mouse splenocytes were surface stained with CoraLite® Plus 488 Anti-Mouse CD3, and 0.25 ug CoraLite® 594 Anti-Mouse TCR beta (H57-597) Armenian Hamster IgG2a RecAb (CL594-65678, Clone: H57-597) or CoraLite® 594 Armenian Hamster IgG Isotype Control (PIP) (CL594-65210, Clone: PIP). Cells were not fixed.



1x10⁶ mouse splenocytes were surface stained with 0.25 ug CoraLite® 594 Anti-Mouse TCR beta (H57-597) Armenian Hamster IgG2a RecAb (CL594-65678, Clone: H57-597) (red) or CoraLite® 594 Armenian Hamster IgG Isotype Control (PIP) (CL594-65210, Clone: PIP) (blue). Cells were not fixed.