For Research Use Only

CoraLite®594-conjugated ATP5A1 Monoclonal antibody

www.ptglab.com

Catalog Number: CL594-66037

Featured Product

Basic Information

Catalog Number: GenBank Accession Number:

CL594-66037 BC064562 GeneID (NCBI):

100ul, Concentration: 1000 ug/ml by 498

Nanodrop: **UNIPROT ID:** P25705 Mouse Full Name:

Isotype ATP synthase, H+ transporting, lgG2b mitochondrial F1 complex, alpha

subunit 1, cardiac muscle Immunogen Catalog Number:

Calculated MW: AG8119

> 60 kDa Observed MW: 50 kDa

Purification Method:

Protein A purification

CloneNo.: 1B10H3

Recommended Dilutions:

WB 1:500-1:1000 IF/ICC 1:50-1:500

Excitation/Emission maxima

wavelengths: 588 nm / 604 nm

Applications

Tested Applications:

WB. IF/ICC

Species Specificity:

human, mouse, rat, monkey

Positive Controls:

WB: HEK-293 cells, HeLa cells, MCF-7 cells, NIH/3T3

IF/ICC: HepG2 cells,

Background Information

The ATP5A1 gene encodes the a subunit of mitochondrial ATP synthase which produces ATP from ADP in the presence of a proton gradient across the membrane. The mitochondrial ATP synthase, also known as Complex V or F1F0 ATP synthase, is a multi-subunit enzyme complex consisting of two functional domains, the F1-containing the catalytic core and the Fo-containing the membrane proton channel. FO domain has 10 subunits: a, b, c, d, e, f, g, OSCP, A6L, and F6. F1 is composed of subunits $\alpha, \beta, \gamma, \delta, \epsilon$, and a loosely attached inhibitor protein IF1. Recently defect in ATP5A1 has been linked to the fatal neonatal mitochondrial encephalopathy. ATP5A1 is localized in the mitochondria and anti-ATP5A1 can be used as the loading control for mitochondrial or Complex V proteins. This antibody recognizes the endogenous ATP5A1 protein in lysates from various cell lines and tissues. The predicted MW of ATP5A1 is 60 kDa, while it undergoes the transit peptide cleavage to become a mature form around 50-55 kDa. Several isoforms of ATP5A1 exist due to the alternative splicing.


Storage

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.


PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using Coralite®594 ATP5A1 antibody (CL594-66037, Clone: 1B10H3) at dilution of 1:100.



Various lysates were subjected to SDS PAGE followed by western blot with CL594-66037 (ATP5A1 antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.