For Research Use Only

## CoraLite®594-conjugated HDAC1 Monoclonal antibody

Catalog Number: CL594-66085



**Purification Method:** 

CloneNo.:

wavelengths: 588 nm / 604 nm

4C4G7

Protein G purification

Recommended Dilutions:

Excitation/Emission maxima

IF/ICC 1:4000-1:16000

**Basic Information** 

Catalog Number: GenBank Accession Number:

CL594-66085 BC000301 GeneID (NCBI):

Nanodrop: **UNIPROT ID:** Q13547 Mouse Full Name:

100ul, Concentration: 1000 ug/ml by 3065

Isotype: histone deacetylase 1 lgG1 Calculated MW:

Immunogen Catalog Number: 55 kDa AG19128 Observed MW: 60-65 kDa

**Applications** 

**Tested Applications:** IF/ICC, FC (Intra)

Species Specificity: human, mouse, rat

Positive Controls:

IF/ICC: A431 cells, HeLa cells

## **Background Information**

Histone deacetylases(HDAC) are a class of enzymes that remove the acetyl groups from the lysine residues leading to the formation of a condensed and transcriptionally silenced chromatin. The protein encoded by this gene belongs to the histone deacetylase/acuc/apha family and is a component of the histone deacetylase complex, which is  $responsible for gene\ expression\ silencing.\ It\ also\ plays\ an\ important\ role\ in\ the\ control\ of\ cell\ proliferation\ and$ differentiation by interacting with RB, p53 and other transcription factors. At least 4 classes of HDAC were identified. As a class I HDAC, HDAC 1 was primarily found in the nucleus. This antibody is raised against residues near the C terminus of human HDAC1. The calculated molecular weight of HDAC1 is 55 kDa, but modified HDAC1 is about 60-65 kDa.

Storage

Storage:


Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer


PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage


## Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using CL594-66085 (HDAC1 antibody) at dilution of 1:100.



1X10^6 HeLa cells were intracellularly stained with 0.2 ug CoraLite®594 Anti-Human HDAC1 (CL594-66085, Clone:4C4G7) (red), or 0.2 ug Control Antibody. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).



Immunofluorescent analysis of (4% PFA) fixed A431 cells using Coralite®594 HDAC1 antibody (CL594-66085, Clone: 4C4G7) at dilution of 1:8000.



Immunofluorescent analysis of (4% PFA) fixed A431 cells using CoraLite®594 HDAC1 antibody (CL594-66085, Clone: 4C4G7) at dilution of 1:8000.