For Research Use Only

## CoraLite®594-conjugated ORM1/2 Monoclonal antibody

Catalog Number: CL594-66097




Basic Information	Catalog Number: CL594-66097	GenBank Accession Number: BC026238	Purification Method: Protein G purification				
	Size: 100ul, Concentration: 1000 ug/ml by Nanodrop; Source: Mouse Isotype: IgG1 Immunogen Catalog Number: AG19248	GeneID (NCBI):	CloneNo.: 5F2H7 Recommended Dilutions: IF/ICC 1:50-1:500 Excitation/Emission maxima wavelengths: 588 nm / 604 nm				
				Applications	Tested Applications: IF/ICC, FC (Intra)	Positive Controls: IF/ICC : HepG2 cells,	
					Species Specificity: human		
				Background Information	Alpha-1-acid glycoprotein 1 (AGP1), also called orosomucoid-1 (ORM1), is a glycoprotein synthesized mostly by hepatocytes and present in human plasma. ORM1 is an acute-phase reactant protein controlled by glucocorticoids, interleukin-1 and interleukin-6, and increase up to 5-50 times upon infection and/or inflammation. Anti-apoptotic effect and role as immunomodulator of ORM have been reported. ORM is an important carrier for synthetic drugs ar influences their distribution and availability in the body. This antibody recognizes a band about 44 kDa in human plasma which may be due to the glycosylation of ORM1 or the dimer formation of the protein. This antibody recognizes both ORM1 and ORM2.		
Storage	Storage: Store at -20°C. Avoid exposure to ligh Storage Buffer:		nent.				
	PBS with 50% Glycerol, 0.05% Proclir	1300, 0.5% BSA, pH 7.3.					


For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll freeE: proteintech@ptglab.comin USA), or 1(312) 455-8498 (outside USA)W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using CL594-66097 (ORM1/2 antibody) at dilution of 1:50.



1X10^6 HepG2 cells were intracellularly stained with 0.4 ug Coralite®594 Anti-Human ORM1/2 (CL594-66097, Clone:5F2H7) (red), or 0.4 ug Mouse IgG1 Isotype Control (CL594-66360, Clone: T1F8D3F10) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).