For Research Use Only

CoraLite®594-conjugated AK2 Monoclonal antibody

Catalog Number:CL594-66127 Featured Product




Basic Information	Catalog Number: CL594-66127	GenBank Accession Number: BC009405	Purification Method: Protein A purification				
	Size: 100ul , Concentration: 1000 ug/ml by Nanodrop; Source: Mouse Isotype: IgG2a Immunogen Catalog Number: AG17911	GeneID (NCBI): 204	CloneNo.: 3H2C10				
		UNIPROT ID: P54819 Full Name: adenylate kinase 2 Calculated MW: 26 kDa Observed MW: 29 kDa	Recommended Dilutions: IF/ICC 1:50-1:500 Excitation/Emission maxima wavelengths: 588 nm / 604 nm				
				Applications	Tested Applications: IF/ICC, FC (Intra)	Positive Controls: IF/ICC : HepG2 cells,	
					Species Specificity: human	IF/ICC : nepuz cetts,	
Background Information	AK2(Adenylate kinase 2, mitochondrial) is also named as ADK2 and belongs to the adenylate kinase family. It plays a unique role in energy metabolism and energy transfer by regulating the ATP/ADP rate between the cytoplasma matrix and the mitochondria and it is increased expression during adipocyte and B cell differentiation(PMID:23020757). AK2 catalyzes the reversible transfer of a phosphoryl group from ATP to AMP(PMID:19043417). Defects in AK2 are the cause of reticular dysgenesis (RDYS)(PMID:19043416). It has 6 isoforms produced by alternative splicing.						
Storage	Storage: Store at -20°C. Avoid exposure to light. Stable for one year after shipment. Storage Buffer: PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3. Aliguoting is unnecessary for -20°C storage						


For technical support and original validation data for this product please contact: T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free E: proteintech@ptglab.com in USA), or 1(312) 455-8498 (outside USA) W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using CL594-66127 (AK2 antibody) at dilution of 1:100.



1X10^6 HepG2 cells were intracellularly stained with 0.4 ug CoraLite® 594 Anti-Human AK2 (CL594-66127, Clone:3H2C 10) (red), or 0.4 ug Mouse IgG2a Isotype Control (CL594-66360-2, Clone: K11A1B2A2) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).