

For Research Use Only

CoraLite®594-conjugated MPO Monoclonal antibody

Catalog Number:CL594-66177

Featured Product

1 Publications



Basic Information

Catalog Number:

CL594-66177

Size:

100ul , Concentration: 1000 ug/ml by

Nanodrop;

Source:

Mouse

Isotype:

IgA

Immunogen Catalog Number:

AG17564

GenBank Accession Number:

BC130476

GeneID (NCBI):

4353

UNIPROT ID:

P05164

Full Name:

myeloperoxidase

Calculated MW:

745 aa, 84 kDa

Observed MW:

90 kDa

Purification Method:

Thiophilic affinity chromatograph

CloneNo.:

4C11F6

Recommended Dilutions:

IF-P 1:50-1:500

Excitation/Emission maxima
wavelengths:

588 nm / 604 nm

Applications

Tested Applications:

IF-P, FC (Intra)

Cited Applications:

IF

Species Specificity:

human, rat

Cited Species:

human

Positive Controls:

IF-P : human appendicitis tissue, human liver tissue

Background Information

The MPO gene encodes myeloperoxidase, a lysosomal hemoprotein located in the azurophilic granules of polymorphonuclear (PMN) leukocytes and monocytes. In response to stimulation, MPO is activated into a transient intermediate with potent antimicrobial oxidizing abilities(PMID:17650507). The mRNA is translated into a single protein of 90 kDa, which displays enzymatic activity and undergoes proteolytic maturation into a heavy chain of 59 kDa and a light chain of 13.5 kDa; these subunits then dimerize into the mature tetramer and the mature MPO is a heterotetramer composed of two identical heavy chains and two identical light chains(PMID:12773517). The 24-kDa material had a map identical to that of 13.5 kDa subunit and represents a dimer of the 13.5 kDa subunit (PMID:3008892). Defects in MPO are the cause of myeloperoxidase deficiency (MPOD). It has 3 isoforms produced by alternative splicing.

Notable Publications

Author	Pubmed ID	Journal	Application
Xing Liu	37734390	Thromb Haemost	IF

Storage

Storage:

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer:

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



1X10⁶ HL-60 cells were intracellularly stained with 0.8 ug CoraLite®594 Anti-Human MPO (CL594-66177, Clone:4C11F6) (red), or 0.8 ug Isotype Control. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).



Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded human appendicitis tissue using CoraLite®594 MPO antibody (CL594-66177, Clone: 4C11F6) at dilution of 1:200. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded human appendicitis tissue using CoraLite®594 MPO antibody (CL594-66177, Clone: 4C11F6) at dilution of 1:200. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).