For Research Use Only

CoraLite®594-conjugated JNK Monoclonal antibody Catalog Number:CL594-66210 Featured Product





Basic Information	Catalog Number: CL594-66210	GenBank Accession Number: BC 130572	Purification Method: Protein G purification				
	Size: 100ul , Concentration: 1000 ug/ml by Nanodrop; Source: Mouse Isotype: IgG1 Immunogen Catalog Number: AG21426	GeneID (NCBI): 5599 UNIPROT ID: P45983 Full Name: mitogen-activated protein kinase 8 Calculated MW: 48 kDa	CloneNo.: 1A12E1 Recommended Dilutions: IF/ICC 1:50-1:500 Excitation/Emission maxima wavelengths: 588 nm / 604 nm				
				Observed MW: 44-48 kDa, 50-55 kDa			
				Applications	Tested Applications:	ICC, FC (Intra) IF/ICC : Jurkat cells, ecies Specificity:	
					Species Specificity: human, mouse, rat		
		Background Information		MAPK8(Mitogen-activated protein kinase 8) is also named as JNK1, PRKM8, SAPK1, SAPK1C and belongs to the MAP kinase subfamily. MAPK8 is activated by dual phosphorylation at a Thr-Pro-Tyr motif during response to UV light. MAPK8 functions to phosphorylate c-Jun at N-terminal serine regulatory sites of Ser-63 and Ser-73, mapping within the transactivation domain. Phosphorylation of these sites in response to UV results in transcriptional activation of c-Jun. It has some isoforms produced by alternative splicing with the molecular weight of 46 kDa and 48 kDa. This protein can be phosphorylated and this antibody recognizes the 46 kDa and 55 kDa bands in western blot(PMID:11062067).			
Storage	Storage: Store at -20°C. Avoid exposure to ligh Storage Buffer: PBS with 50% Glycerol, 0.05% Proclin Aliquoting is unnecessary for -20°C s	n300, 0.5% BSA, pH 7.3.					

For technical support and original validation data for this product please contact: T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free E: proteintech@ptglab.com in USA), or 1(312) 455-8498 (outside USA) W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data





 1X10^6 HeLa cells were intracellularly stained with 0.8 ug CoraLite®594 Anti-Human JNK (CL594-66210, Clone:1A12E1) (red), or 0.8 ug CoraLite®594 Mouse IgG1 Isotype Control (MOPC-21) (CL594-65124, Clone: MOPC-21) (blue). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).
 Immunofluores Jurkat cells usin (CL594-66210, MOPC-21), CL594-65124, Clone: MOPC-21) (blue). Cells were fixed and permeabilized with Transcription Factor

Immunofluorescent analysis of (4% PFA) fixed Jurkat cells using CoraLite®594 JNK antibody (CL594-66210, Clone: 1A12E1) at dilution of 1:200.