### For Research Use Only

# CoraLite®594-conjugated Beta Tubulin Monoclonal antibody

proteintech®
Antibodies | ELISA kits | Proteins
www.ptglab.com

**Purification Method:** 

Protein A purification

Catalog Number:CL594-66240 3 Publications

**Basic Information** 

Catalog Number: GenBank Accession Number:

CL594-66240 BC000748
Size: GeneID (NCBI):

Size:GeneID (NCBI):CloneNo.:100ul , Concentration: 1000 ug/ml by 103811D4A4

Nanodrop; UNIPROT ID: Recommended Dilutions:
Source: Q13509 IF/ICC 1:50-1:500
Mouse Full Name of Source States (Finishing Page)

Mouse Full Name: Excitation/Emission maxima Isotype: tubulin, beta 3 wavelengths:

IgG2a Calculated MW: 588 nm / 604 nm

IgG2a Calculated MW:
Immunogen Catalog Number: 450 aa, 50 kDa
AG0117 Channel MW:

0117 Observed MW: 50-55 kDa

**Applications** 

Tested Applications: IF/ICC, FC (Intra) Cited Applications:

--

IF

Species Specificity:

human, mouse, rat, pig, zebrafish, nematode

Positive Controls:

IF/ICC : SH-SY5Y cells,

#### **Background Information**

There are five tubulins in human cells: alpha, beta, gamma, delta, and epsilon. Tubulins are conserved across species. They form heterodimers, which multimerize to form a microtubule filament. An alpha and beta tubulin heterodimer is the basic structural unit of microtubules. The heterodimer does not come apart, once formed. The alpha and beta tubulins, which are each about 55 kDa MW, are homologous but not identical. Alpha and beta tubulins have been widely used as loading controls. Tubulin expression may vary according to resistance to antimicrobial and antimitotic drugs.

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Sun K Kim	39314338	bioRxiv	IF
Wen Lu	37722034	Proc Natl Acad Sci U S A	IF
Wen Lu	36945460	bioRxiv	IF

Storage

Storage:


Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffe


PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

## Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed SH-SY5Y cells using the CoraLite®594-conjugated version of this antibody, CL594-66240 (beta Tubulin antibody), at dilution of 1:100.



1X10^6 HeLa cells were intracellularly stained with 0.4 ug CoraLite®594 Anti-Human Beta Tubulin (CL594-66240, Clone:1D4A4) (red), or 0.4 ug CoraLite®594 Mouse IgCa Isotype Control (C1.18.4) (CL594-65208, Clone: C1.18.4) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).