For Research Use Only

CoraLite®594-conjugated PKC Iota Monoclonal antibody



Purification Method:

wavelengths:

593 nm / 614 nm

Catalog Number: CL594-66493

Basic Information

Catalog Number: GenBank Accession Number:

CL594-66493 BC022016 Protein A purification
Size: Genel D (NCBI): CloneNo.:

100ul , Concentration: 1000 µg/ml by 5584 3A1B6
Nanodrop; Full Name: Recommended Dilutions:

Source: protein kinase C, iota IF 1:50-1:500

Mouse Calculated MW: Excitation/Emission maxima

Isotype: 68 kDa IgG2a

Immunogen Catalog Number:

AG4990

Positive Controls:

IF: HeLa cells,

Applications

Tested Applications: FC (Intra), IF

Species Specificity: Human, rat, mouse

Background Information

The atypical protein kinase C isoform PRKC iota (PRKCI) is a member of the protein kinase C (PKC) family of serine/threonine protein kinases. PKC family comprises at least eight members, which are differentially expressed and are involved in a wide variety of cellular processes. PRKC iota is calcium-independent and phospholipid-dependent. It is not activated by phorbolesters or diacylglycerol. This kinase can be recruited to vesicle tubular clusters (VTCs) by direct interaction with the small GTPase RAB2, where this kinase phosphorylates glyceraldehyde-3-phosphate dehydrogenase (GAPD/GAPDH) and plays a role in microtubule dynamics in the early secretory pathway. This kinase is found to be necessary for BCL-ABL-mediated resistance to drug-induced apoptosis and therefore protects leukemia cells against drug-induced apoptosis. PRKC iota plays a key role in cell proliferation, differentiation, and carcinogenesis, and it has been shown to be a human oncogene.

Storage

Storage:

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.


Storage Buffer

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using CL594-66493 (PKC iota antibody) at dilution of 1:50.



1X10^6 HeLa cells were intracellularly stained with 0.2 ug CoraLite® 594 Anti-Human PKC lota (CL594-66493, Clone:3A1B6) (red), and 0.2 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with 0.1% TritonX-100.