For Research Use Only

CoraLite®594-conjugated PARP1 Monoclonal antibody

www.ptglab.com

Catalog Number:CL594-66520

Featured Product

Basic Information

Catalog Number: GenBank Accession Number:

CL594-66520 BC037545 GeneID (NCBI):

100ul, Concentration: 1000 ug/ml by 142 Nanodrop: **UNIPROT ID:** P09874

Mouse Full Name:

Isotype: poly (ADP-ribose) polymerase 1 lgG1

Calculated MW: Immunogen Catalog Number: 1014 aa, 113 kDa AG19173 Observed MW:

113-116 kDa, 85-89 kDa

Purification Method:

Protein G purification

CloneNo.: 1D7D4

Recommended Dilutions: IF/ICC 1:50-1:500

Excitation/Emission maxima

wavelengths: 588 nm / 604 nm

Applications

Tested Applications: IF/ICC, FC (Intra)

Species Specificity: human, mouse, rat

Positive Controls:

IF/ICC: HeLa cells, Neuro-2a cells

Background Information

PARP1 (poly(ADP-ribose) polymerase 1) is a nuclear enzyme catalyzing the poly(ADP-ribosyl)ation of many key proteins in vivo. The normal function of PARP1 is the routine repair of DNA damage. Activated by DNA strand breaks, the PARP1 is cleaved into an 85 to 89-kDa COOH-terminal fragment and a 24-kDa NH2-terminal peptide by caspases during the apoptotic process. The appearance of PARP fragments is commonly considered as an important biomarker of apoptosis. In addition to caspases, other proteases like calpains, cathepsins, granzymes and matrix metalloproteinases (MMPs) have also been reported to cleave PARP1 and gave rise to fragments ranging from 42-89-kD. This antibody was generated against the N-terminal region of human PARP1 and it recognizes the full-length as well as the cleavage of the PARP1.

Storage

Storage:


Store at -20°C. Avoid exposure to light. Stable for one year after shipment. Storage Buffer

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.


Aliquoting is unnecessary for -20°C storage

in USA), or 1(312) 455-8498 (outside USA)


Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using CL594-66520 (PARP1 antibody) at dilution of 1:100.



1X10^6 HeLa cells were intracellularly stained with 0.4 ug CoraLite®594 Anti-Human PARP1 (CL594-66520, Clone:1D7D4) (red), or 0.4 ug Control Antibody. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).



Immunofluorescent analysis of (4% PFA) fixed Neuro-2a cells using Coralite®594 PARP1 antibody (CL594-66520, Clone: 1D7D4) at dilution of 1:2000, CL488-Phalloidin (green).