For Research Use Only

## CoraLite®594-conjugated NXN Monoclonal antibody

Catalog Number: CL594-66603



**Purification Method:** 

Protein G purification

Recommended Dilutions:

Excitation/Emission maxima

IF/ICC 1:50-1:500

wavelengths: 588 nm / 604 nm

CloneNo.:

2C6A4

**Basic Information** 

Catalog Number: GenBank Accession Number:

nucleoredoxin

CL594-66603 BC009327

GeneID (NCBI): 100ul, Concentration: 1000 ug/ml by 64359

Nanodrop: **UNIPROT ID:** Q6DKJ4

Mouse Full Name:

lgG1 Calculated MW: Immunogen Catalog Number: 435 aa, 48 kDa

AG9248 Observed MW: 48 kDa

Positive Controls:

IF/ICC: HEK-293 cells,

**Applications** 

**Tested Applications:** IF/ICC, FC (Intra) Species Specificity: human, mouse, rat

Isotype:

## **Background Information**

Nucleoredoxin (NRX/NXN), a relatively uncharacterized member of the TRX family protein, has recently been reported to regulate the Wnt/ $\beta$ -catenin pathway, which itself regulates cell fate and early development, in a redoxdependent manner (PMID:17567240). Northern blot analysis detected several nucleoredoxin splice variants in adult mouse tissues, with highest expression in testis and skin and lowest expression in liver(PMID:9119370). It has 3 isoforms produced by alternative splicing. This antibody is specific to NXN.


Storage

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.


PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

## Selected Validation Data



Immunofluorescent analysis of (-20°C Ethanol) fixed HEK-293 cells using CoraLite® 594 NXN antibody (CL594-66603, Clone: 2C6A4) at dilution of 1:200.



1X10^6 HeLa cells were intracellularly stained with 0.4 ug CoraLite® 594 Anti-Human NXN (CL594-66603, Clone:2C6A4) (red), or 0.4 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).