For Research Use Only

## CoraLite®594-conjugated BAG6 Monoclonal antibody Catalog Number:CL594-66661 Featured Product





Basic Information	Catalog Number: CL594-66661	GenBank Accession Number: BC003133	Purification Method: Protein G purification				
	Size: 100ul , Concentration: 1000 ug/ml by Nanodrop; Source: Mouse Isotype: IgG1 Immunogen Catalog Number: AG24525	GeneID (NCBI):	CloneNo.: 1B8D3 Recommended Dilutions: WB 1:500-1:2000 Excitation/Emission maxima wavelengths: 588 nm / 604 nm				
				Applications	Tested Applications: WB, FC (Intra)	Positive Controls: WB : HeLa cells, LNCaP cells, HepG2 cells	
					Species Specificity: human, mouse, rat		
				Background Information	BAT3 also known as Scythe or BAG6, is a nuclear protein implicated in the control of apoptosis and natural killer (NK) cell-dendritic cell (DC) interaction. BAT3 was first discovered as a member of a group of genes located within the class III region of the human major histocompatibility complex on chromosome 6, and has been extensively studied for its role in regulating apoptosis under various stress conditions such as DNA damage and endoplasmic reticulum-related stress. BAT3 has been shown to be required for p53 acetylation, which is critical for the enhancement of p53 transcriptional activity in response to DNA damage. In addition, BAT3 is involved in the regulation of development and reproduction of mammals by acting as a co-chaperone of the heat shock protein HSP70.		
					HSP70.		

For technical support and original validation data for this product please contact: T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free E: proteintech@ptglab.com in USA), or 1(312) 455-8498 (outside USA) W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data





Various lysates were subjected to SDS PAGE followed by western blot with CL594-66661 (BAG6 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.

1X10^6 HeLa cells were intracellularly stained with 0.4 ug CoraLite®594 Anti-Human BAG6 (CL594-66661, Clone:1B8D3) (red), or 0.4 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).