For Research Use Only

CoraLite®594-conjugated Phospho-RIPK1 (Ser161) Monoclonal antibody

Catalog Number: CL594-66854





Basic Information	Catalog Number: CL594-66854	GenBank Accession Number: NM_003804	Purification Method: Protein G purification
	Size: 100ul , Concentration: 1000 ug/ml by Nanodrop; Source: Mouse Isotype: IgG1	GenelD (NCBI): 8737	CloneNo.: 1B2G1
		UNIPROT ID: Q13546 Full Name: receptor (TNFRSF)-interacting serine- threonine kinase 1 Calculated MW: 76 kDa Observed MW: 74 kDa, 45 kDa	Recommended Dilutions: IF/ICC 1:50-1:500 Excitation/Emission maxima wavelengths: 588 nm / 604 nm
Applications	Tested Applications: IF/ICC, FC (Intra) Species Specificity: human	Positive Controls: IF/ICC : Calyculin A treated HeLa cells,	
Background Information	RIPK1(Receptor-interacting serine/threonine-protein kinase 1) is primarily involved in mediating TNF-R1-induced cell activation, apoptosis and necroptosis and belongs to a novel class of kinases thatfunction in cell survival and cell death mechanisms(PMID:22685397). It has 2 isoforms produced by alternative splicing. It also can exist as the protein with lower molecular weight about 25 kDa and 45 kDa in HAEC and HUVEC (PMID: 22685397).		
Storage	Storage: Store at -20°C. Avoid exposure to light. Stable for one year after shipment. Storage Buffer: PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3. Aliquoting is unnecessary for -20°C storage		

For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)E: proteintech@ptglab.comW: ptglab.comW: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data





Immunofluorescent analysis of (4% PFA) fixed Calyculin A treated HeLa cells using Coralite®594 Phospho-RIPK1 (Ser161) antibody (CL594-66854, Clone: 1B2G1) at dilution of 1:200.

1X10^6 HEK-293T cells untreated (dashed lines) or treated with Calyculin A (red) were intracellularly stained with 0.13 ug Coralite®594 Anti-Human Phospho-RIPK1 (Ser161) (CL594-66854, Clone:1B2C1) (red), or 0.13 ug Control Antibody (blue). Cells were fixed with 4% PFA and permeabilized with 90% MeOH.