For Research Use Only

CoraLite®594-conjugated mTOR Monoclonal antibody



Purification Method:

Catalog Number: CL594-66888

Featured Product

2 Publications

Basic Information

Catalog Number: GenBank Accession Number: CL594-66888 NM_004958

2L594-66888 NM_004958 Protein A purification dize: GeneID (NCBI): CloneNo.:

Size: GeneID (NCBI): CloneNo. 100ul , Concentration: 1000 ug/ml by 2475 1G11A3

Nanodrop; UNIPROT ID: Recommended Dilutions:
Source: P42345 IF/ICC 1:50-1:500

Mouse Full Name: Excitation/Emission maxima

Isotype: FK506 binding protein 12-rapamycin wavelengths:

 IgG2a
 associated protein 1
 588 nm / 604 nm

 Immunogen Catalog Number:
 Calculated MW:

Immunogen Catalog Number:Calculated MWAG28395289 kDaObserved MW:

250-289 kDa

Applications

Tested Applications:

IF/ICC

Cited Applications:

IF

Species Specificity: Human, Mouse, Rat Cited Species:

human, mouse

Positive Controls:

IF/ICC : HepG2 cells,

Background Information

MTOR, also named as FRAP1, FRAP, FRAP2 and RAPT1, belongs to the PI3/PI4-kinase family. MTOR is a Ser/Thr protein kinase that functions as an ATP and amino acid sensor to balance nutrient availability and cell growth. MTOR is kinase subunit of both mTORC1 and mTORC2, which regulate cell growth and survival in response to nutrient and hormonal signals. mTORC1 is activated in response to growth factors or amino-acids. mTORC2 is also activated by growth factors, but seems to be nutrient-insensitive. mTORC2 seems to function upstream of Rho GTPases to regulate the actin cytoskeleton, probably by activating one or more Rho-type guanine nucleotide exchange factors. mTORC2 promotes the serum-induced formation of stress-fibers or F-actin. MTOR has a calculated molecular mass of 289 kDa, and always can be detected at about 250 kDa due to some modifications (PMID: 14578359).

Notable Publications

Author	Pubmed ID	Journal	Application
Shuo Geng	37961551	bioRxiv	IF
Margot R F Reijnders	37057673	Genet Med	IF

Storage

Storage


Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using CoraLite®594 mTOR antibody (CL594-66888, Clone: 1G11A3) at dilution of 1:200.