For Research Use Only

CoraLite®594-conjugated FBL Monoclonal antibody

Catalog Number: CL594-66985



Purification Method:

Protein G purification

Recommended Dilutions:

Excitation/Emission maxima

IF/ICC 1:50-1:500

wavelengths: 588 nm / 604 nm

CloneNo.:

3A9E8

Basic Information

Catalog Number: GenBank Accession Number:

CL594-66985 BC019260
Size: GeneID (NCBI):

100ul, Concentration: 1000 ug/ml by 2091 Nanodrop:

Nanodrop; UNIPROT ID: 5ource: P22087

Mouse Full Name: Isotype: fibrillarin

IgG1 Calculated MW:
Immunogen Catalog Number: 321 aa, 34 kDa
AG8841 Observed MW:

Observe 34 kDa

Applications
Tested Applications:

IF/ICC, FC (Intra)
Species Specificity:

human, mouse, rat

Positive Controls:

IF/ICC : HepG2 cells,

Background Information

Fibrillarin (Fbl) is an essential nucleolar protein that plays an essential role in ribosome biogenesis and more particularly in the methylation of ribosomal RNAs and rDNA histones. FBL is highly expressed in various cancers and correlates with poor survival outcomes in cancer patients, such as hepatocellular carcinoma (PMID: 33376525). Knockdown of FBL sensitizes tumor cells and xenografts to DNA crosslinking agents, and leads to homologous recombination-mediated DNA repair defects (PMID: 37489617).

Storage

Storage:


Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer:


PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using CoraLite®594 FBL antibody (CL594-66985, Clone: 3A9E8) at dilution of 1:200.



1x10^6 HepG2 cells were intracellularly stained with 0.8 ug Coralite®594 Anti-Human FBL (CL594-66985, Clone:3A9E8)(red), or 0.8 ug Coralite®594 Mouse IgG1 Isotype Control (MOPC-21) (CL594-65124, Clone: MOPC-21) (blue). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).