For Research Use Only

CoraLite®594-conjugated TAK1 Monoclonal antibody



Purification Method:

IF/ICC 1:50-1:500

wavelengths: 588 nm / 604 nm

Excitation/Emission maxima

CloneNo.:

Protein G purification

Catalog Number: CL594-67707

Basic Information

Catalog Number: GenBank Accession Number:

CL594-67707 BC017715 GeneID (NCBI):

100ul , Concentration: 1000 $\mu g/ml$ by 6885 1A5B2 Nanodrop: **UNIPROT ID:** Recommended Dilutions:

043318 Mouse Full Name:

Isotype: mitogen-activated protein kinase lgG1 kinase kinase 7

Immunogen Catalog Number: Calculated MW: AG21286 579 aa. 64 kDa

> Observed MW: 67-70 kDa

Applications

Tested Applications:

Species Specificity: Human, Mouse, Rat Positive Controls:

IF/ICC: HepG2 cells,

Background Information

MAP3K7(Mitogen-activated protein kinase kinase kinase 7) is also named as TAK1 and belongs to the MAP kinase kinase kinase subfamily. It plays an important role in the cascades of cellular responses evoked by changes in the environment. It has been linked to interleukin-1 receptor and tumor necrosis factor receptor signaling (PMID: 16186825). It has 4 isoforms (53-55 kDa, 64-70 kDa and 75-80 kDa) produced by alternative splicing.


Storage

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using CoraLite®594 TAK1 antibody (CL594-67707, Clone: 1A5B2) at dilution of 1:200.