For Research Use Only

CoraLite®594-conjugated HADHB Monoclonal antibody

IgG2b

www.ptglab.com

Purification Method:

588 nm / 604 nm

Catalog Number: CL594-67967

Basic Information

Catalog Number: GenBank Accession Number: BC017564

CL594-67967 Protein A purification GeneID (NCBI): CloneNo.:

100ul, Concentration: 1000 ug/ml by 3032 1D12F4 Nanodrop: **UNIPROT ID:** Recommended Dilutions:

P55084 IF/ICC 1:50-1:500 Mouse

Full Name: Excitation/Emission maxima Isotype hydroxyacyl-Coenzyme A wavelengths: dehydrogenase/3-ketoacyl-

Coenzyme A thiolase/enoyl-Immunogen Catalog Number:

Coenzyme A hydratase (trifunctional AG30315

protein), beta subunit

Calculated MW:

51 kDa

Observed MW: 47 kDa

Applications

Tested Applications: IF/ICC, FC (Intra) Species Specificity:

human, mouse, rat, pig

Positive Controls:

IF/ICC: HeLa cells,

Background Information

HADHB, also named as TP- beta, Acetyl-CoA acyltransferase and Beta-ketothiolase, is a mitochondrial trifunctional enzyme subunit beta. Mitochondrial trifunctional enzyme catalyzes the last three of the four reactions of the mitochondrial beta-oxidation pathway. The mitochondrial beta-oxidation pathway is the major energy-producing process in tissues and is performed through four consecutive reactions breaking down fatty acids into acetyl-CoA. Among the enzymes involved in this pathway, the trifunctional enzyme exhibits specificity for long-chain fatty acids. Mitochondrial trifunctional enzyme is a heterotetrameric complex composed of two proteins, the trifunctional enzyme subunit alpha/HADHA carries the 2,3-enoyl-CoA hydratase and the 3-hydroxyacyl-CoA dehydrogenase activities, while the trifunctional enzyme subunit beta/HADHB described here bears the 3-ketoacyl-CoA thiolase activity. HADHB has 2 isoforms produced by alternative splicing with the MW of 49 kDa and 51 kDa.

Storage

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.


Aliquoting is unnecessary for -20°C storage

in USA), or 1(312) 455-8498 (outside USA)

Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using Coralite®594 HADHB antibody (CL594-67967, Clone: 1D12F4) at dilution of 1:200.



1X10^6 HeLa cells were intracellularly stained with 0.4 ug CoraLite® 594 Anti-Human HADHB (CL594-67967, Clone:1D12F4) (red), or 0.4 ug Mouse IgG2b Isotype Control (CL594-66360-3, Clone: K11B8C4B5) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).