For Research Use Only

## CoraLite®594-conjugated MYBBP1A Monoclonal antibody

www.ptglab.com

Catalog Number: CL594-67996

**Basic Information** 

Catalog Number: GenBank Accession Number:

CL594-67996 BC050546 GeneID (NCBI):

100ul, Concentration: 1000 ug/ml by 10514 Nanodrop: **UNIPROT ID:** Q9BQG0 Mouse Full Name:

Isotype: MYB binding protein (P160) 1a

lgG1 Calculated MW: Immunogen Catalog Number: 149 kDa

AG6008 Observed MW: 140 kDa

**Applications** 

**Tested Applications:** IF/ICC, FC (Intra)

Species Specificity: Human, mouse, rat

**Purification Method:** 

Protein G purification

CloneNo.: 1H2D10

Positive Controls:

IF/ICC: HepG2 cells,

Recommended Dilutions: IF/ICC 1:50-1:500

Excitation/Emission maxima

wavelengths: 588 nm / 604 nm

**Background Information** 

 $The protoon cogene \ MYB is predominantly \ expressed in immature \ hemopoietic \ cells \ where \ it \ has \ an \ essential \ role \ in \ decreases \ for \ in \ decreases \$ hemopoietic cell proliferation and differentiation. Oncogenically activated forms of MYB is generally N- and/or Cterminal truncations of the normal MYB protein. Removal of the C terminus of MYB disrupts or deletes a region termed the negative regulatory domain (NRD), resulting in an increase in DNA binding, transactivation, and transformation by MYB. One feature of the NRD is a leucine zipper-like motif [PMID: 8302594]. Murine Myb-binding protein-1a (MYBBP1A), originally called P160, was identified by its ability to interact specifically with Myb via this leucine zipper-like motif. MYBBP1A modulates MYB activity upon binding to the MYB NRD [PMID: 10644447, 9447996].

Storage


Storage:

Store at -20°C. Avoid exposure to light. Stable for one year after shipment. Storage Buffer


PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

## Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using CoraLite® 594 MYBBP1A antibody (CL594-67996, Clone: 1H2D10) at dilution of 1:200.



1X10^6 HepG2 cells were intracellularly stained with 0.4 ug Coralite®594 Anti-Human MYBBP1A (CL594-67996, Clone:1H2D10) (red), or 0.4 ug Control Antibody. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).