For Research Use Only

CoraLite®594-conjugated MYBBP1A Monoclonal antibody

proteintech®
Antibodies | ELISA kits | Proteins
www.ptglab.com

Purification Method:

wavelengths:

588 nm / 604 nm

Catalog Number: CL594-67996

Basic Information

Catalog Number: GenBank Accession Number:

CL594-67996 BC050546 Protein G purification
Size: Genel D (NCBI): CloneNo.:

100ul , Concentration: 1000 µg/ml by 10514 1H2D10

Nanodrop; Full Name: Recommended Dilutions:
Source: MYB binding protein (P160) 1a IF 1:50-1:500

Source: MYB binding protein (P160) 1a IF 1:50-1:500

Mouse Calculated MW: Excitation/Emission maxima

Isotype: 149 kDa IgG1 Observed MW:

Immunogen Catalog Number: 140 kDa

AG6008

Applications

Tested Applications:

IF

Species Specificity: Human, mouse, rat **Positive Controls:**

IF : HepG2 cells,

Background Information

The protooncogene MYB is predominantly expressed in immature hemopoietic cells where it has an essential role in hemopoietic cell proliferation and differentiation. Oncogenically activated forms of MYB is generally N- and/or C-terminal truncations of the normal MYB protein. Removal of the C terminus of MYB disrupts or deletes a region termed the negative regulatory domain (NRD), resulting in an increase in DNA binding, transactivation, and transformation by MYB. One feature of the NRD is a leucine zipper-like motif [PMID: 8302594]. Murine Myb-binding protein-1a (MYBBP1A), originally called P160, was identified by its ability to interact specifically with Myb via this leucine zipper-like motif. MYBBP1A modulates MYB activity upon binding to the MYB NRD [PMID: 106444447, 9447996].

Storage

Storage:

Store at -20°C. Avoid exposure to light.


Storage Buffer:

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using CoraLite® 594 MYBBP1A antibody (CL594-67996, Clone: 1H2D10) at dilution of 1:200.



1X10^6 HepG2 cells were intracellularly stained with 0.4 ug Coralite®594 Anti-Human MYBBP1A (CL594-67996, Clone:1H2D10) (red), or 0.4 ug Control Antibody. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).