For Research Use Only

CoraLite®594-conjugated DYKDDDDK tag Recombinant antibody (Binds to FLAG® tag epitope) Catalog Number:CL594-80801-2




Basic Information	Catalog Number: CL594-80801-2 Size:	GenBank Accession Number: GeneID (NCBI): 8 Full Name: Flag Tag	Purification Method: Protein A purification CloneNo.:
	100ul , Concentration: 1000 µg/ml by Nanodrop;		240568A11 Recommended Dilutions:
	Source: Rabbit Isotype: IgG		IF/ICC: 1:50-1:500 Excitation/Emission maxima
			wavelengths: 588 nm / 604 nm
	Immunogen Catalog Number: AG2329		
Applications	Tested Applications: IF/ICC		Controls:
	Species Specificity: recombinant protein	IF/ICC : Transfected HEK-293 cells,	
Background Information	Protein tags are protein or peptide sequences located either on the C- or N- terminal of the target protein, which facilitates one or several of the following characteristics: solubility, detection, purification, localization and expression. The DYKDDDDK(FLAG) peptide has been used extensively as a general tag in expression vectors. This peptide can be expressed and detected with the protein of interest as an amino-terminal or carboxy-terminal fusion. N-terminal DDDDK vectors provide an Ek cleavage site for removal of the fusion tag. The DDDDK peptide is likely to be located on the surface of a fusion protein because of its hydrophilic nature. As a result, the DDDDK peptide is more likely to be accessible to antibodies. A DDDDK-tag can be used in many different assays that require recognition by an antibody, such as western blotting, immunocytochemistry, immunoprecipitation, flow cytometry, protein purification, and in the study of protein-protein interactions, cell ultrastructure, and protein localization and so on. This antibody is a rabbit recombinant antibody raised against 3xFlag (3xDYKDDDDKT) sequence and recognizes the (1x) and (3x)DYKDDDDK peptide and detects DDDDK-tagged proteins. Anti-FLAG is a registered trademark of Sigma-Aldrich Biotechnology.		
Storage	Storage: Store at -20°C. Avoid exposure to ligh Storage Buffer: PBS with 50% glycerol, 0.05% Proclin Aliquoting is unnecessary for -20°C s	1300, 0.5% BSA, pH7.3	ient.

For technical support and original validation data for this product please contact: T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free E: proteintech@ptglab.com in USA), or 1(312) 455-8498 (outside USA) W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Immunofluorescent analysis of (-20°C Ethanol) fixed Transfected HEK-293 cells using CoraLite®594 DYKDDDDK tag antibody (CL594-80801-2, Clone: 240568A11) at dilution of 1:200.