For Research Use Only

CoraLite®594-conjugated Phospho-NFкВ p65 (Ser468) Recombinant antibody

Catalog Number: CL594-82335


-	
	proteintech®
jy –	Antibodies ELISA kits Proteins
	www.ptglab.com

Basic Information	Catalog Number: CL594-82335 Size:	GenBank Accession Number: BC011603 GeneID (NCBI):	Purification Method: Protein A purification CloneNo.:	
	100ul , Concentration: 1000 ug/ml by Nanodrop; Source: Rabbit Isotype: IgG	5970 UNIPROT ID: Q04206 Full Name: v-rel reticuloendotheliosis viral oncogene homolog A (avian) Calculated MW: 65 kDa Observed MW: 75 kDa	6N1 Excitation/Emission maxima wavelengths: 588 nm / 604 nm	
Applications	Tested Applications: FC (Intra) Species Specificity: human, mouse			
Background Information	Nuclear factor kB (NF-kB) is a collective term for a small family of dimeric transcription factors [comprising p65 (RelA) and RelB, c-Rel, p50/p105 (NF-kB1), and p52/p100 (NF-kB2)]. All NF-kB proteins share a Rel homology domain (RHD), which is responsible for DNA binding and dimerization. Only p65, RelB, and c-Rel contain potent transactivation domains within sequences from the C-terminal to the RHD. Exterior signals lead to the phosphorylation and degradation of the inhibitory complex IkB, which is modulated by the IkB kinase (IKK), and its degradation allows for the release of the typical NF-kB heterodimer, p65/p50, to translocate into the nucleus. NF-kB binds to its cognate DNA elements and can transcriptionally activate different target genes among which 200-500 genes have been implicated in cell survival/apoptosis, cell growth, immune response, and inflammation.			
Storage	Storage: Store at -20°C. Avoid exposure to ligh Storage Buffer: PBS with 50% glycerol, 0.05% Proclin Aliquoting is unnecessary for -20°C s	1300, 0.5% BSA, pH7.3		

For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll freeE: proteintech@ptglab.comin USA), or 1(312) 455-8498 (outside USA)W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



1X10^6 NIH/3T3 cells untreated (dashed lines) or treated with Calyculin A which intracellularly stained with 0.13 ug Coralite®594 Phospho-NF-Kb P65 (Ser468) Recombinant Antibody (CL594-82335, Clone:6N1)(red), or 0.13 ug Coralite®594 Rabbit IgG Isotype Control RecAb (CL594-98136, Clone: 240953C9) (blue). Cells were fixed with 4% PFA and permeabilized with 90% MeOH.