For Research Use Only

CoraLite®594 Anti-Mouse CD79a Rabbit Recombinant Antibody

Catalog Number: CL594-98129



Purification Method:

Protein A purification

Excitation/Emission maxima

wavelengths:

588 nm / 604 nm

Basic Information

Catalog Number: GenBank Accession Number:

CL594-98129 NM_007655.3

 Size:
 GeneID (NCBI):
 CloneNo.:

 100ug, 500 ug/ml
 12518
 241231F7

Source: UNIPROT ID: Recommended Dilutions:

Rabbit P11911 FC (Intra): 0.25 ug per 10^6 cells in Isotype: Full Name: 100 µl suspension

Isotype: Full Name:

IgG CD79A antigen (immunoglobulin-

Immunogen Catalog Number: associated alpha)

EG1388 Calculated MW:

25 kDa

Positive Controls:

FC (Intra): mouse splenocytes,

Applications

Tested Applications:

FC (Intra)

Species Specificity:

mouse

Background Information

CD79a, also named as B-cell antigen receptor complex-associated protein alpha chain or MB-1 membrane glycoprotein, is a 226 amino acid protein, which contains one ITAM domain and one Ig-like C2-type (immunoglobulin-like) domain. CD79A is expressed in B cell and localizes in the cell membrane. CD79A is required in cooperation with CD79B for initiation of the signal transduction cascade activated by binding of antigen to the B-cell antigen receptor complex (BCR) which leads to internalization of the complex, trafficking to late endosomes and antigen presentation. CD79A is also required for BCR surface expression and for efficient differentiation of pro-and pre-B-cells.

Storage


Storage:

Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.


Storage Buffer

PBS with 0.09% sodium azide, pH7.3

Selected Validation Data



1x10^6 mouse splenocytes were intracellularly stained with 0.25 ug CoraLite®594 Anti-Mouse CD79a Rabbit RecAb (CL594-98129, Clone:241231F7) or 0.25 ug CoraLite®594 Rabbit IgG Isotype Control RecAb (CL594-98136, Clone: 240953C9). Cells were co-stained with CoraLite® Plus 647 Anti-Mouse CD45R (B220) (RA3-6B2) (CL647-65139, Clone: RA3-6B2). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).



1x10^6 mouse splenocytes were intracellularly stained with 0.25 ug Coralite®594 Anti-Mouse CD79a Rabbit RecAb (CL594-98129, Clone: 241231F7)(red) or Coralite®594 Rabbit IgG Isotype Control RecAb (CL594-98136, Clone: 240953C9) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).