For Research Use Only

CoraLite® Plus 647-conjugated Podoplanin Polyclonal antibody



Catalog Number: CL647-11629

Basic Information

Catalog Number: GenBank Accession Number:

BC022812

43 kDa

CL647-11629 GeneID (NCBI):

100ul, Concentration: 1000 µg/ml by 10630

Full Name: Source: podoplanin Rabbit Calculated MW: Isotype: 17 kDa, 25 kDa IgG Observed MW: Immunogen Catalog Number:

Purification Method: Antigen affinity purification Excitation/Emission maxima

wavelengths: 654 nm / 674 nm

Applications

Tested Applications:

FC (Intra)

Species Specificity:

human

Background Information

Podoplanin was identified as a 43 kDa glycoprotein found in the cell membranes of glomerular epithelial cells (podocyte) (PMID: 9327748). It is a lymphatic marker because the expression of podoplanin has been detected in lymphatic but not blood vascular endothelium, and is useful as the marker of tumor-associated Lymphangiogenesis. Podoplanin has a function in developing testis, most likely at the level of cell-cell interactions among pre-meiotic germ cells and immature Sertoli cells. It may be involved in cell migration and/or actin cytoskeleton organization. When expressed in keratinocytes, PDPN induces changes in cell morphology with transfected cells showing an elongated shape, numerous membrane protrusions, major reorganization of the actin cytoskeleton, increased motility and decreased cell adhesion. It is required for normal lung cell proliferation and alveolus formation at birth. PDPN induces platelet aggregation. It does not have any effect on folic acid or amino acid transport and does not function as a water channel or as a regulator of aquaporin-type water channels. The antibody is specific to Podoplanin.

Storage

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.


Storage Buffer:

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

Selected Validation Data



1X10^6 HEK-293 cells were intracellularly stained with 0.2 ug Coralite® Plus 647 Anti-Human Podoplanin (CL647-11629) (red), or 0.2 ug Control Antibody (blue). (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).