For Research Use Only

CoraLite® Plus 647-conjugated iNOS Polyclonal antibody

Catalog Number:CL647-18985

Featured Product

1 Publications





Basic Information	Catalog Number: CL647-18985	GenBank Accession Nur NM_000625	mber: Purification Method: Antigen affinity purification
	Size: 100ul , Concentration: 1000 µg/ml by	GenelD (NCBI): 4843	Recommended Dilutions: IF/ICC 1:50-1:500
	Nanodrop; Source: Rabbit Isotype: IgG	UNIPROT ID: Excitation/Emission maxima P35228 wavelengths: Full Name: 654 nm / 674 nm nitric oxide synthase 2, inducible Calculated MW: 131 kDa Observed MW: 110-130 kDa, 65-70 kDa	
Applications	Tested Applications: IF/ICC, FC (Intra) Species Specificity: human, mouse, rat Cited Species: mouse		Positive Controls: IF/ICC : HepG2 cells,
Background Information	NOS2, also named as iNOS and NOS2A, produces nitric oxide (NO) which is a messenger molecule with diverse functions throughout the body. NO is a reactive free radical which acts as a biologic mediator in several processes, including neurotransmission, antimicrobial and antitumoral activities. NOS2 is a nitric oxide synthase which is expressed in liver and is inducible by a combination of lipopolysaccharide and certain cytokines. iNOS has a very short half-life due to rapid degradation by calpain. iNOS monomer is a direct substrate of calpain I and can be cleaved by calpain I at the canonical CaM-binding site(503-532aa) of iNOS, and then a ~70kD band can be detected by western (PMID:11786228). This antibody is specific to NOS2.		
Notable Publications	Author Pub	med ID Journal	l Application
	Fengan Cao 398	06403 Respir F	Res
Storage	Storage: Store at -20°C. Avoid exposure to ligh Storage Buffer: PBS with 50% Glycerol, 0.05% Proclin Aliquoting is unnecessary for -20°C st	n300, 0.5% BSA, pH 7.3.	ter shipment.

For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll freeE: proteintech@ptglab.comin USA), or 1(312) 455-8498 (outside USA)W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data





1X10⁶ HepG2 cells were intracellularly stained with 0.2 ug CoraLite® Plus 647 Anti-Human iNOS (CL647-18985) (red), or 0.2 ug isotype control (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).

Immunofluorescent analysis of (-20°C Methanol) fixed HepG2 cells using CoraLite® Plus 647 iNOS antibody (CL647-18985) at dilution of 1:100.