## For Research Use Only

## CoraLite® Plus 647-conjugated PI3 Kinase p110 Beta Polyclonal antibody




Catalog Number:CL647-20584 Featured Product			
<b>Basic Information</b>	Catalog Number: CL647-20584	GenBank Accession Number: NM_006219	Purification Method: Antigen affinity purification
	Size: 100ul , Concentration: 1000 µg/ml by Nanodrop; Source: Rabbit Isotype: IgG	GeneID (NCBI): / 5291 UNIPROT ID: P42338 Full Name: phosphoinositide-3-kinase, catalytic beta polypeptide Calculated MW: 123 kDa Observed MW: 110-130 kDa	Excitation/Emission maxima wavelengths: 654 nm / 674 nm
Applications	Tested Applications: FC (Intra) Species Specificity: human, rat		
Background Information	PIK3CB(phosphatidylinositol 4,5-bisphosphate 3-kinase catalytic subunit beta isoform) is also named as PIK3C1, PI3K-beta, p110beta. The gene encodes a 1070 amino acid protein which belongs to the PI3/PI4-kinase family. Phosphoinositide 3-kinases (PI3Ks) have been implicated as participants in signaling pathways regulating cell growth by virtue of their activation in response to various mitogenic stimuli. The class I PI3 kinases are heterodimers composed of 110 kDa catalytic subunits that associate with regulatory adaptor proteins. Four class I catalytic subunits have been identified, PIK3CA (p110α), PIK3CB (p110β), PIK3CD (p110δ) and PIK3CG (p110γ) (PMID:19177002). This antibody is specific to PIK3CB.		
Storage	Storage: Store at -20°C. Avoid exposure to light. Stable for one year after shipment. Storage Buffer: PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3. Aliquoting is unnecessary for -20°C storage		

For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free<br/>in USA), or 1(312) 455-8498 (outside USA)E: proteintech@ptglab.comW: ptglab.comW: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data



1X10^6 HeLa cells were intracellularly stained with 0.2 ug CoraLite® Plus 647 Anti-Human Pl3 Kinase p110 Beta (CL647-20584) (red), or 0.2 ug isotype control (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).