For Research Use Only

## CoraLite® Plus 647-conjugated MTCO2 Polyclonal antibody

www.ptglab.com

**Purification Method:** 

IF/ICC 1:50-1:500

wavelengths: 654 nm / 674 nm

Antigen affinity purification

Excitation/Emission maxima

Recommended Dilutions:

Catalog Number: CL647-55070

Featured Product

**Basic Information** 

Catalog Number: GenBank Accession Number:

CL647-55070 YP\_003024029 GeneID (NCBI):

100ul , Concentration: 1000  $\mu g/ml$  by 4513 Nanodrop: Full Name:

Source: cytochrome c oxidase II

Rabbit Calculated MW:

Isotype: 26 kDa

IgG Observed MW:

23-26 kDa

**Applications** 

**Tested Applications:** IF/ICC, FC (Intra)

Species Specificity: human, mouse, rat

Positive Controls:

IF/ICC: HepG2 cells,

## **Background Information**

COX2, also named as COII, COXII and MTCO2, belongs to the cytochrome c oxidase subunit 2 family. It is the component of the respiratory chain that catalyzes the reduction of oxygen to water. Subunits 1-3 form the functional core of the enzyme complex. Subunit 2 transfers the electrons from cytochrome c via its binuclear copper A center to the bimetallic center of the catalytic subunit 1. Defects in COX2 are a cause of mitochondrial complex IV deficiency (MT-C4D). The antibody is specific to COX2.

Storage

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.


PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

## Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using CoraLite® Plus 647 MTCO2 antibody (CL647-55070) at dilution of 1:200.



1x10^6 HeLa cells were intracellularly stained with 0.8 ug CoraLite® Plus 647-conjugated MTCO2 Polyclonal antibody (CL647-55070)(red), or 0.8 ug CoraLite® Plus 647-conjugated Rabbit 1gG control Rabbit PolyAb (CL647-30000) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).