## CoraLite® Plus 647-conjugated PPAR **proteintech** Antibodies | ELISA kits | Proteins Gamma Monoclonal antibody www.ptglab.com

Catalog Number:CL647-60127

## 44 **Basic Info**

Basic Information	Catalog Number: CL647-60127	GenBank Accession Number: BC006811	Purification Method: Protein G purification	
	Size: 100ul , Concentration: 1000 µg/ml by Nanodrop; Source: Mouse Isotype: IgG1 Immunogen Catalog Number: AG10005	GenelD (NCBI): 5468	CloneNo.: 4E12F10	
		UNIPROT ID: P37231 Full Name: peroxisome proliferator-activated receptor gamma	Excitation/Emission maxima wavelengths: 654 nm / 674 nm	
				Calculated MW: 58 kDa
		Observed MW: 50-60 kDa	MM:	
		Applications	Tested Applications: Species Specificity: human, mouse, rat	
Background Information	Peroxisome Proliferator-Activated Receptors (PPARs) are ligand-activated intracellular transcription factors, members of the nuclear hormone receptor superfamily (NR), that includes estrogen, thyroid hormone receptors, retinoic acid, Vitamin D3 as well as retinoid X receptors (RXRs). The PPAR subfamily consists of three subtypes encoded by distinct genes denoted PPARa (NR1C1), PPARβ/ō (NR1C2) and PPARy (NR1C3), which are activated by selective ligands. PPARy, also named as PPARG, contains one nuclear receptor DNA-binding domain and is a receptor that binds peroxisome proliferators such as hypolipidemic drugs and fatty acids. It plays an important role in the regulation of lipid homeostasis, adipogenesis, INS resistance, and development of various organs. Defects in PPARG are the cause of familial partial lipodystrophy type 3 (FPLD3) and may be associated with susceptibility to obesity. Defects in PPARG can lead to type 2 INS-resistant diabetes and hypertension. PPARG mutations may be associated with colon cancer. Genetic variations in PPARG are associated with susceptibility to glioma type 1 (GLM1). PPARG has two isoforms with molecular weight 57 kDa and 54 kDa (PMID: 9831621), but modified PPARG is about 67 KDa (PMID: 16809887). PPARG2 is a splice variant and has an additional 30 amino acids at the N-terminus (PMID: 15689403). Experimental data indicate that a 45 kDa protein displaying three different sequences immunologically related to the nuclear receptor PPARG2 is located in mitochondria (mt-PPAR). However, the molecular weight of this protein is clearly less when compared to that of PPARG2 (57 kDa). (PMID: 10922459). PPARG has been reported to be localized mainly (but not always) in the nucleus. PPARG can also be detected in the cytoplasm and was reported to possess extra-nuclear/non-genomic actions (PMID: 17611413; 19432669; 14681322).			
Storage	Storage: Store at -20°C. Avoid exposure to light. Stable for one year after shipment. Storage Buffer. PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3. Aliquoting is unnecessary for -20°C storage			

For technical support and original validation data for this product please contact: T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free E: proteintech@ptglab.com in USA), or 1(312) 455-8498 (outside USA) W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data