For Research Use Only

CoraLite® Plus 647 Anti-Human M-CSF (3G12E6)




Catalog Number: CL647-65283


Basic Information	Catalog Number: CL647-65283	GenBank Accession Number: BC021117	Purification Method: Affinity purification
	Size: 100ug , 0.2 mg/mL	GenelD (NCBI): 1435	CloneNo.: 3G12E6
	Source: Mouse Isotype: IgG1 Immunogen Catalog Number: HZ-1192	Full Name: colony stimulating factor 1 (macrophage) Calculated MW: 60 kDa	Excitation/Emission maxima wavelengths: 654 nm / 674 nm
	Species Specificity: Human		
Background Information	M-CSF plays an essential role in the regulation of survival, proliferation and differentiation of hematopoietic precursor cells, especially mononuclear phagocytes. M-CSF promotes the release of proinflammatory chemokines, and thereby plays an important role in innate immunity and in inflammatory processes. It also plays an important role in the regulation of osteoclast proliferation and differentiation, the regulation of bone resorption, and is required for normal bone development. On the cellular level, M-CSF promotes reorganization of the actin cytoskeleton, regulates formation of membrane ruffles, cell adhesion and cell migration and plays a role in lipoprotein clearance (PMID: 29734839; 29679908; 28202039)		
Storage	Storage: Store at 2-8°C. Avoid exposure to lig Storage Buffer: PBS with 0.09% sodium azide and 0	yht. Stable for one year after shipment. .5% BSA.	

For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)E: proteintech@ptglab.comW: ptglab.comW: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data





1X10^{^6} LPS and TNF-a then Brefeldin A treated human PBMCs were surface stained with Coralite® Plus 555 Anti-Human CD11b then intracellularly stained with 0.2 ug Coralite® Plus 647 Anti-Human M-CSF (CL647-65283, Clone:3G12E6) or Mouse IgG1 Isotype Control. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011). Monocytes were gated.

1X10^6 LPS and TNF-a then Brefeldin A treated human PBMCs were intracellularly stained with 0.2 ug CoraLite® Plus 647 Anti-Human M-CSF (CL647-65283, Clone:3G12E6) (red) or Mouse IgG1 Isotype Control (blue), or unactivated then Brefeldin A treated human PBMCs were intracellularly stained with 0.2 ug CoraLite® Plus 647 Anti-Human M-CSF (CL647-65283, Clone:3G12E6) (black, dashed line). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit