For Research Use Only

CoraLite® Plus 647 Anti-Human IL-10 (7B8G3)



Catalog Number: CL647-65286

Basic Information

Catalog Number:

CL647-65286

Size: 100ug , 0.2 mg/mL

Source: Mouse Isotype:

lgG2b 21 kDa

Immunogen Catalog Number:

HZ-1145

Purification Method:

Affinity purification CloneNo.:

7B8G3

Excitation/Emission maxima

wavelengths: 654 nm / 674 nm

Applications

Tested Applications:

FC (Intra)

Species Specificity:

Human

Background Information

Interleukin (IL)-10 is an anti-inflammatory cytokine, produced by T helper (Th) cells, macrophages, monocytes, and B cells, that plays a crucial role in preventing inflammatory and autoimmune pathologies. It downregulates the expression of Th1 cytokines, MHC class II antigens, and co-stimulatory molecules on macrophages. It also enhances $B\ cell\ survival,\ proliferation,\ and\ antibody\ production.\ IL-10\ can\ block\ NF-\kappa B\ activity,\ and\ is\ involved\ in\ the$ regulation of the JAK-STAT signaling pathway. IL-10, along with its receptors, describes an important role in pathogenesis of various diseases, including infectious, inflammatory, autoimmune diseases. IL-10 mutations are associated with an increased susceptibility to HIV-1 infection and rheumatoid arthritis.

GenBank Accession Number:

BC104252

Full Name:

interleukin 10

Calculated MW:

3586

GeneID (NCBI):


Storage

Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.


Storage Buffer:

PBS with 0.09% sodium azide and 0.5% BSA.

Selected Validation Data



1X10^6 LPS and Brefeldin A treated human PBMCs were intracellularly stained with 0.2 ug CoraLite® Plus 647 Anti-Human IL-10 (CL647-65286, Clone:7B8G3) or Mouse IgG2b Isotype Control. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011). Monocytes were gated.



1X10^6 LPS and Brefeldin A treated human PBMCs were intracellularly stained with 0.2 ug Coralite® Plus 647 Anti-Human IL-10 (CL647-65286, Clone:7B8G3) (red) or Mouse IgG2b Isotype Control (blue), or unactivated and Brefeldin A treated human PBMCs were intracellularly stained with 0.2 ug Coralite® Plus 647 Anti-Human IL-10 (CL647-65286, Clone:7B8G3) (black, dashed line). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).

