## For Research Use Only

## CoraLite® Plus 647 Anti-Human CD8a (OKT8) Mouse IgG2a Recombinant Antibody



Catalog Number: CL647-65604

**Basic Information** 

Catalog Number:

CL647-65604

Size:

100tests, 5 ul/test

Source: Mouse Isotype:

IgG2a

GenBank Accession Number:

BC025715 GeneID (NCBI):

925

**ENSEMBL Gene ID:** ENSG00000153563

UNIPROT ID: P01732

Full Name: CD8a molecule

Calculated MW: 235 aa, 26 kDa

**Purification Method:** 

Protein A purification

CloneNo.: ОКТ8

**Recommended Dilutions:** 

FC: 5 ul per 10^6 cells in a 100 µl

suspension

Excitation/Emission maxima

wavelengths: 654 nm / 674 nm

**Applications** 

**Tested Applications:** 

Species Specificity:

human

**Positive Controls:** 

FC: human PBMCs,

## **Background Information**

CD8 is a transmembrane glycoprotein composed of two disulfide-linked chains. It can be present as a homodimer of CD8a or as a heterodimer of CD8a and CD8β (PMID: 3264320; 8253791). CD8 is found on most thymocytes. The majority of class I-restricted T cells express mostly the CD8 $\alpha\beta$  heterodimer while CD8 $\alpha\alpha$  homodimers alone have been found on some gut intraepithelial T cells, on some T cell receptor (TCR)  $\gamma \delta$  T cells and on NK cells (PMID: 2111591; 1831127; 8420975). CD8 acts as a co-receptor that binds to MHC class-I and participates in cytotoxic T-cell activation (PMID: 8499079). During T cell development, CD8 is required for positive selection of CD4-/CD8+ T cells (PMID: 1968084).


Storage

Storage:


Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.

PBS with 0.09% sodium azide and 0.5% BSA, pH7.3

## **Selected Validation Data**



1x10^6 human PBMCs were surface stained with PE Anti-Human CD3 (OKT3) Mouse IgG2a Recombinant Antibody (PE-65569, Clone: OKT3), and 5 ul Coralite® Plus 647 Anti-Human CD8a (OKT8) Mouse IgG2a RecAb (CL647-65604, Clone: OKT8) or Coralite® Plus 647 Mouse IgG2a Isotype Control (C1.18.4) (CL647-65208, Clone: C1.18.4). Cells were not fixed.



1x10^6 human PBMCs were surface stained with 5 ul CoraLite® Plus 647 Anti-Human CD8a (OKT8) Mouse IgG2a RecAb (CL647-65604, Clone:OKT8) (red), or CoraLite® Plus 647 Mouse IgG2a Isotype Control (C1.18.4) (CL647-65208, Clone: C1.18.4) (blue). Cells were not fixed.