

For Research Use Only

CoraLite® Plus 647-conjugated HDAC1 Monoclonal antibody



Catalog Number:CL647-66085

Basic Information

Catalog Number: CL647-66085	GenBank Accession Number: BC000301	Purification Method: Protein G purification
Size: 100ul , Concentration: 1000 µg/ml by Nanodrop;	GeneID (NCBI): 3065	CloneNo.: 4C4G7
Source: Mouse	Full Name: histone deacetylase 1	Recommended Dilutions: IF 1:500-1:2000
Isotype: IgG1	Calculated MW: 55 kDa	Excitation/Emission maxima wavelengths: 650 nm / 665 nm
Immunogen Catalog Number: AG19128	Observed MW: 65 kDa	

Applications

Tested Applications: FC (Intra), IF	Positive Controls: IF : A431 cells,
Species Specificity: human, mouse, rat	

Background Information

Histone deacetylases(HDAC) are a class of enzymes that remove the acetyl groups from the lysine residues leading to the formation of a condensed and transcriptionally silenced chromatin. The protein encoded by this gene belongs to the histone deacetylase/acuc/apha family and is a component of the histone deacetylase complex, which is responsible for gene expression silencing. It also plays an important role in the control of cell proliferation and differentiation by interacting with RB, p53 and other transcription factors. At least 4 classes of HDAC were identified. As a class I HDAC, HDAC 1 was primarily found in the nucleus. This antibody is raised against residues near the C terminus of human HDAC1. The calculated molecular weight of HDAC1 is 55 kDa, but modified HDAC1 is about 60-65 kDa .

Storage


Storage:
Store at -20°C. Avoid exposure to light.
Storage Buffer:
PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA) E: proteintech@ptglab.com W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed A431 cells using CoraLite® Plus 647 HDAC1 antibody (CL647-66085, Clone: 4C4G7) at dilution of 1:1000.



1X10⁶ HeLa cells were intracellularly stained with 0.4 ug CoraLite® Plus 647 Anti-Human HDAC1 (CL647-66085, Clone:4C4G7) (red), or 0.4 ug Control Antibody. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).