For Research Use Only

CoraLite® Plus 647-conjugated DEK Monoclonal antibody



Catalog Number: CL647-66194

Featured Product

Basic Information

Catalog Number: GenBank Accession Number:

CL647-66194 BC035259 GeneID (NCBI):

100ul , Concentration: 1000 $\mu g/mL$ by 7913 Nanodrop:

UNIPROT ID: P35659 Mouse Full Name: Isotype: **DEK** oncogene IgG2a Calculated MW: Immunogen Catalog Number: 43 kDa

AG8225 Observed MW:

40-50 kDa

Purification Method:

Protein A purification

CloneNo.: 5E3A6

Excitation/Emission maxima

wavelengths: 654 nm / 674 nm

Applications

Tested Applications:

FC (Intra)

Species Specificity: human, mouse, rat

Background Information

Dek oncogene(DEK) is first recognized in a fusion with CAN nucleoporin protein, and it's a also autoantigen in patients with pquciarticular onset juvenile rheumatoid arthritis. As a site-specific binding protein, DEK may involve in transcriptional regulation and signal transdcution. Analysis by agarose gel celctrophoresis, DEK showed to reduce the efficiency with which chromation is replicated. It's one of the first components of the splicing complex, which remains associate with spliced exons dependent on prior splicing of pre-mRNA.


Storage

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



1x10^6 HeLa cells were intracellularly stained with 0.4 ug CoraLite® Plus 647-conjugated DEK Monoclonal antibody (CL647-66194, Clone:5E3A6) (red), or 0.4 ug CoraLite® Plus 647 Mouse IgG2a Isotype Control (C1.18.4) (CL647-65208, Clone: C1.18.4) (blue). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).